

13 Topology of \mathbb{R} II

14 Topology of \mathbb{R} III

15 Topology of \mathbb{R} IV

16 Topology of \mathbb{R} V

17 Topology of \mathbb{R} VI

Topology of \mathbb{R}

Intervals



Open interval:

$$(a, b) = \{x : a < x < b\}$$


Closed interval:

$$[c, d] = \{x : c \leq x \leq d\}$$

Half-open interval:

$$(e, f] = \{x : e < x \leq f\}$$

Interior point




Definition (Interior point)

If $E \subseteq \mathbb{R}$ then x is an **interior point** of E if x lies in an open interval that is contained in E , i.e., $\exists c > 0$ such that $(x - c, x + c) \subset E$.

Interior point examples

Set E	Interior points?
$(-1, 1)$	Every point
$[0, 1]$	Every point <i>except the endpoints</i>
\mathbb{N}	\nexists
\mathbb{R}	Every point
\mathbb{Q}	\nexists
$(-1, 1) \cup [0, 1]$	Every point <i>except 1</i>
$(-1, 1) \setminus \{\frac{1}{2}\}$	Every point


Neighbourhood



Definition (Neighbourhood)

A **neighbourhood** of a point $x \in \mathbb{R}$ is an open interval containing x .

Deleted neighbourhood




Definition (Deleted neighbourhood)

A **deleted neighbourhood** of a point $x \in \mathbb{R}$ is a set formed by removing x from a neighbourhood of x .

$$(a, b) \setminus \{x\}$$

Isolated point



$$E = (a_1, b_1) \cup [a_2, b_2) \cup \{x\}$$

Definition (Isolated point)

If $x \in E \subseteq \mathbb{R}$ then x is an **isolated point** of E if there is a neighbourhood of x for which the only point in E is x itself, *i.e.*, $\exists c > 0$ such that $(x - c, x + c) \cap E = \{x\}$.

Isolated point examples

Set E	Isolated points?
$(-1, 1)$	\nexists
$[0, 1]$	\nexists
\mathbb{N}	Every point
\mathbb{R}	\nexists
\mathbb{Q}	\nexists
$(-1, 1) \cup [0, 1]$	\nexists
$(-1, 1) \setminus \{\frac{1}{2}\}$	\nexists



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn


Lecture 13
Topology of \mathbb{R}^n
Wednesday 4 October 2017

Last time...

Concepts associated with sets of real numbers:

- Countable set
- Interval
- Neighbourhood
- Deleted neighbourhood
- Interior point
- Isolated point

Accumulation point



$$E = \left\{ 1 - \frac{1}{n} : n \in \mathbb{N} \right\}$$

Definition (Accumulation Point or Limit Point)

If $E \subseteq \mathbb{R}$ then x is an **accumulation point** or **limit point** of E if every neighbourhood of x contains infinitely many points of E ,

$$\text{i.e.,} \quad \forall c > 0 \quad (x - c, x + c) \cap (E \setminus \{x\}) \neq \emptyset.$$

Notes:

- It is possible but not necessary that $x \in E$.
- The shorthand condition is equivalent to saying that every deleted neighbourhood of x contains at least one point of E .

Accumulation point examples

Set E	Accumulation points?
$(-1, 1)$	$[-1, 1]$
$[0, 1]$	$[0, 1]$
\mathbb{N}	\nexists
\mathbb{R}	\mathbb{R}
\mathbb{Q}	\mathbb{R}
$(-1, 1) \cup [0, 1]$	$[-1, 1]$
$(-1, 1) \setminus \{\frac{1}{2}\}$	$[-1, 1]$
$\{1 - \frac{1}{n} : n \in \mathbb{N}\}$	$\{1\}$

Boundary point



Definition (Boundary Point)

If $E \subseteq \mathbb{R}$ then x is a **boundary point** of E if every neighbourhood of x contains at least one point of E and at least one point not in E , i.e.,

$$\begin{aligned} \forall c > 0 \quad & (x - c, x + c) \cap E \neq \emptyset \\ & \wedge (x - c, x + c) \cap (\mathbb{R} \setminus E) \neq \emptyset. \end{aligned}$$

Note: It is possible but not necessary that $x \in E$.

Definition (Boundary)

If $E \subseteq \mathbb{R}$ then the **boundary** of E , denoted ∂E , is the set of all boundary points of E .

Boundary point examples

Set E	Boundary points?
$(-1, 1)$	$\{-1, 1\}$
$[0, 1]$	$\{0, 1\}$
\mathbb{N}	\mathbb{N}
\mathbb{R}	\nexists
\mathbb{Q}	\mathbb{R}
$(-1, 1) \cup [0, 1]$	$\{-1, 1\}$
$(-1, 1) \setminus \{\frac{1}{2}\}$	$\{-1, \frac{1}{2}, 1\}$
$\{1 - \frac{1}{n} : n \in \mathbb{N}\}$	$\{1 - \frac{1}{n} : n \in \mathbb{N}\} \cup \{1\}$

Closed set



Definition (Closed set)

A set $E \subseteq \mathbb{R}$ is **closed** if it contains all of its accumulation points.

Definition (Closure of a set)

If $E \subseteq \mathbb{R}$ and E' is the set of accumulation points of E then $\overline{E} = E \cup E'$ is the **closure** of E .

Note: If the set E has no accumulation points, then E is closed because there are no accumulation points to check.

Open set



Definition (Open set)

A set $E \subseteq \mathbb{R}$ is **open** if every point of E is an **interior point**.

Definition (Interior of a set)

If $E \subseteq \mathbb{R}$ then the **interior** of E , denoted E° or E° , is the set of all **interior points** of E .

Examples

Set E	Closed?	Open?	\bar{E}	E°	∂E
$(-1, 1)$	NO	YES	$[-1, 1]$	E	$\{-1, 1\}$
$[0, 1]$	YES	NO	E	$(0, 1)$	$\{0, 1\}$
\mathbb{N}	YES	NO	\mathbb{N}	\emptyset	\mathbb{N}
\mathbb{R}	YES	YES	\mathbb{R}	\mathbb{R}	\emptyset
\emptyset					
\mathbb{Q}					
$(-1, 1) \cup [0, 1]$					
$(-1, 1) \setminus \{\frac{1}{2}\}$					
$\{1 - \frac{1}{n} : n \in \mathbb{N}\}$					



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 14
Topology of \mathbb{R}^n III
Friday 6 October 2017

Announcements

- **Assignment 3** is due on Friday 20 Oct 2017 at 4:25pm
(remember cover sheet!)
- **Math 3A03 Test #1**
Monday 23 Oct 2017 at 7:00pm in MDCL 1102

Examples

Set E	Closed?	Open?	\bar{E}	E°	∂E
$(-1, 1)$	NO	YES	$[-1, 1]$	E	$\{-1, 1\}$
$[0, 1]$	YES	NO	E	$(0, 1)$	$\{0, 1\}$
\mathbb{N}	YES	NO	\mathbb{N}	\emptyset	\mathbb{N}
\mathbb{R}	YES	YES	\mathbb{R}	\mathbb{R}	\emptyset
\emptyset	YES	YES	\emptyset	\emptyset	\emptyset
\mathbb{Q}	NO	NO	\mathbb{R}	\emptyset	\mathbb{R}
$(-1, 1) \cup [0, 1]$	NO	NO	$[-1, 1]$	$(-1, 1)$	$\{-1, 1\}$
$(-1, 1) \setminus \{\frac{1}{2}\}$	NO	YES	$[-1, 1]$	E	$\{-1, \frac{1}{2}, 1\}$
$\{1 - \frac{1}{n} : n \in \mathbb{N}\}$	NO	NO	$E \cup \{1\}$	\emptyset	$E \cup \{1\}$

Component intervals of open sets

What does the most general open set look like?

Theorem (Component intervals)

If G is an open subset of \mathbb{R} and $G \neq \emptyset$ then there is a unique (possibly finite) sequence of disjoint open intervals $\{(a_n, b_n)\}$ such that

$$G = (a_1, b_1) \cup (a_2, b_2) \cup \cdots \cup (a_n, b_n) \cup \cdots ,$$

$$\text{i.e., } G = \bigcup_{n=1}^{\infty} (a_n, b_n) .$$

*The open intervals (a_n, b_n) are said to be the **component intervals** of G .*

(Textbook (TBB) [Theorem 4.15, p. 231](#))

Component intervals of open sets

Main ideas of proof of [component intervals theorem](#):

- $x \in G \implies x$ is an interior point of $G \implies$
 - some neighbourhood of x is contained in G ,
i.e., $\exists c > 0$ such that $(x - c, x + c) \subseteq G$
 - \exists a largest neighbourhood of x that is contained in G : this largest neighbourhood is $I_x = (\alpha, \beta)$, where

$$\alpha = \inf\{a : (a, x] \subset G\}, \quad \beta = \sup\{b : [x, b) \subset G\}$$

- I_x contains a rational number, *i.e.*, $\exists r \in I_x \cap \mathbb{Q}$
- \therefore We can index all the intervals I_x by rational numbers
- \therefore There are at most countably many intervals that make up G (*i.e.*, G is the union of a sequence of intervals)
- We can choose a disjoint subsequence of these intervals whose union is all of G (see [proof in textbook](#) for details).

Open vs. Closed Sets

Definition (Complement of a set of real numbers)

If $E \subseteq \mathbb{R}$ then the **complement** of E is the set

$$E^c = \{x \in \mathbb{R} : x \notin E\}.$$

Theorem (Open vs. Closed)

If $E \subseteq \mathbb{R}$ then E is open iff E^c is closed.

(Textbook (TBB) [Theorem 4.16](#))

Open vs. Closed Sets

Theorem (Properties of open sets of real numbers)

- 1 The sets \mathbb{R} and \emptyset are open.
- 2 Any *intersection* of a *finite* number of open sets is open.
- 3 Any *union* of an *arbitrary* collection of open sets is open.
- 4 The complement of an open set is closed.

(Textbook (TBB) [Theorem 4.17](#))

Theorem (Properties of closed sets of real numbers)

- 1 The sets \mathbb{R} and \emptyset are closed.
- 2 Any *union* of a *finite* number of closed sets is closed.
- 3 Any *intersection* of an *arbitrary* collection of closed sets is closed.
- 4 The complement of a closed set is open.

(Textbook (TBB) [Theorem 4.18](#))



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 15
Topology of \mathbb{R}^n IV
Monday 16 October 2017

Announcements

- **Assignment 3** is due this Friday 20 Oct 2017 at 4:25pm
(remember cover sheet!)
- **Math 3A03 Test #1, one week from today,**
Monday 23 Oct 2017 at 7:00pm in MDCL 1102
(room is booked for 90 minutes; you should not feel rushed)
- **Math 3A03 Final Exam:** Thurs 21 Dec 2017, 4:00pm–6:30pm

Concepts covered recently

- Countable set
- Interval
- Neighbourhood
- Deleted neighbourhood
- Interior point
- Isolated point
- Accumulation point
- Boundary point
- Boundary
- Closed set
- Closure
- Open set
- Interior
- Complement

Local vs. Global properties

Definition (Bounded function)


A real-valued function f is **bounded** on the set E if there exists $M > 0$ such that $|f(x)| \leq M$ for all $x \in E$.

(i.e., the function f is bounded on E iff $\{f(x) : x \in E\}$ is a bounded set.)


Note: This is a *global* property because there is a single bound M associated with the entire set E .

Example

The function $f(x) = 1/(1 + x^2)$ is bounded on \mathbb{R} . e.g., $M = 1$.




Local vs. Global properties



$f(x) = 1/x$ is not bounded on the interval $E = (0, 1)$.

Local vs. Global properties



$f(x) = 1/x$ is **locally bounded** on the interval $E = (0, 1)$,
i.e., $\forall x \in E, \exists \delta_x, M_x > 0 \mid |f(t)| \leq M_x \forall t \in (x - \delta_x, x + \delta_x)$.

Local vs. Global properties

Definition (Locally bounded at a point)

A real-valued function f is **locally bounded** at the point x if there is a neighbourhood of x in which f is bounded, *i.e.*, there exists $\delta_x > 0$ and $M_x > 0$ such that $|f(t)| \leq M_x$ for all $t \in (x - \delta_x, x + \delta_x)$.

Definition (Locally bounded on a set)

A real-valued function f is **locally bounded** on the set E if f is locally bounded at each point $x \in E$.

Note: The size of the neighbourhood (δ_x) and the local bound (M_x) depend on the point x .

Local vs. Global properties

Example (Function that is not even locally bounded)

Give an example of a function that is defined on the interval $(0, 1)$ but is not locally bounded on $(0, 1)$.

(solution on board)

Example (Function that is a mess near 0)

Give an example of a function $f(x)$ that is defined everywhere, yet in any neighbourhood of the origin there are infinitely many points at which f is not locally bounded.

(solution on board)

Extra Challenge Problem: Is there a function $f : \mathbb{R} \rightarrow \mathbb{R}$ that is not locally bounded anywhere?

Local vs. Global properties

- What condition(s) rule out such pathological behaviour?
- When does a property holding locally (near any given point in a set) imply that it holds globally (for the set as a whole)?
- For example: What condition(s) must a set $E \subseteq \mathbb{R}$ satisfy in order that a function f that is **locally bounded** on E is necessarily **bounded** on E ?
- We will see that the condition we are seeking is that the set E must be “**compact**” ...

Compactness

Recall the Bolzano-Weierstrass theorem, which we proved when investigating sequences of real numbers:

Theorem (Bolzano-Weierstrass theorem for sequences)

Every bounded sequence in \mathbb{R} contains a convergent subsequence.

For any set of real numbers, we define:

Definition (Bolzano-Weierstrass property)

A set $E \subseteq \mathbb{R}$ is said to have the **Bolzano-Weierstrass property** iff any sequence of points chosen from E has a subsequence that converges to a point in E .

Compactness

Theorem (Bolzano-Weierstrass theorem for sets)

A set $E \subseteq \mathbb{R}$ has the *Bolzano-Weierstrass property* iff E is closed and bounded.

(solution on board) (Textbook (TBB) [Theorem 4.21, p. 241](#))

Notes:

- Why do we need both *closed* and *bounded*? Why didn't we need *closed* in the original version of the [Bolzano-Weierstrass theorem](#) (for sequences)?
 - Because we didn't require the limit of the convergent subsequence to be in the set!
- The [Bolzano-Weierstrass theorem for sets](#) implies that "If $E \subseteq \mathbb{R}$ is bounded then its closure \overline{E} has the Bolzano-Weierstrass property".
 - The original [Bolzano-Weierstrass theorem for sequences](#) is a special case of this statement because any convergent sequence together with its limit is a closed set.



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 16
Topology of \mathbb{R}^n
Wednesday 18 October 2017

Announcements

- Niky Hristov's last Monday office hour will be on 23 October, before the midterm test. For the remainder of the term, Niky's office hours will change to Tuesdays 1:30–3:30pm.
- **Assignment 3** is due this Friday 20 Oct 2017 at 4:25pm (remember cover sheet!)
- **Math 3A03 Test #1**,
Monday 23 Oct 2017 at 7:00pm in MDCL 1102
(room is booked for 90 minutes; you should not feel rushed)
 - Question sheets for 2016 tests are now posted on the course wiki.
- **Math 3A03 Final Exam**: Thurs 21 Dec 2017, 4:00pm–6:30pm

Bijections

The terms **one-to-one** (injective), **onto** (surjective), and **one-to-one correspondence** (bijection) are giving some students trouble.

(Recall, we used **bijection** in our definition of **countable**.)

Let's take a step back and recall:

- When we define a **function**, we need three things:
 - the **domain**, *i.e.*, the set to which the function is applied;
 - the **codomain**, *i.e.*, the target set where the values of the function lie;
 - a rule for taking elements of the domain into the codomain.
- If we write $f : A \rightarrow B$ then A is the domain and B is the codomain.
- The **range** of a function is the subset of the codomain consisting of all values of the function applied to the domain.

Bijections

Example

Let $f(x) = x^2$, $x \in \mathbb{R}$.

- Is f onto \mathbb{R} ?
- Is f one-to-one on \mathbb{R} ? On any interval?
- Is f a bijection?

Example

- Find a bijection between $[0, \infty)$ to $[1, \infty)$.
- Find a different bijection between $[0, \infty)$ to $[1, \infty)$.

Extra Challenge Problem:

Construct a bijection between $[0, 1]$ and $(0, 1)$.

Compactness

Definition (Open Cover)

Let $E \subseteq \mathbb{R}$ and let \mathcal{U} be a family of open intervals. If for every $x \in E$ there exists at least one interval $U \in \mathcal{U}$ such that $x \in U$, i.e.,

$$E \subseteq \bigcup \{U : U \in \mathcal{U}\},$$

then \mathcal{U} is called an **open cover** of E .

Example (Open covers of \mathbb{N})

Give examples of open covers of \mathbb{N} .

- $\mathcal{U} = \left\{ \left(n - \frac{1}{2}, n + \frac{1}{2} \right) : n = 1, 2, \dots \right\}$
- $\mathcal{U} = \{(0, \infty)\}$
- $\mathcal{U} = \{(0, \infty), \mathbb{R}, (\pi, 27)\}$

Compactness

Example (Open covers of $\{\frac{1}{n} : n \in \mathbb{N}\}$)

- $\mathcal{U} = \{(0, 1), (0, 2), \mathbb{R}, (\pi, 27)\}$
- $\mathcal{U} = \{(0, 2)\}$
- $\mathcal{U} = \left\{ \left(\frac{1}{n}, \frac{1}{n} + \frac{3}{4} \right) : n = 1, 2, \dots \right\}$

Example (Open covers of $[0, 1]$)

- $\mathcal{U} = \{(-2, 2)\}$
- $\mathcal{U} = \{(-\frac{1}{2}, \frac{1}{2}), (0, 2)\}$
- $\mathcal{U} = \left\{ \left(\frac{1}{n}, 2 \right) : n = 1, 2, \dots \right\} \cup \left\{ \left(-\frac{1}{2}, \frac{1}{2} \right) \right\}$



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 17
Topology of \mathbb{R}^n VI
Friday 20 October 2017

Announcements

- Assignment 3 was due at 4:25pm today.
 - Solutions will be posted tonight.
- Tutorials will occur as usual on Monday, but they will just be Q&A to help with last-minute questions before the test. Tuesday tutorials are CANCELLED this week.
- **Math 3A03 Test #1:**
 - **Monday 23 Oct 2017 at 7:00pm in MDCL 1102**
(room is booked for 90 minutes; you should not feel rushed)
 - Question sheets for 2016 tests are now posted on the course wiki.
 - Bring your student ID, pens, pencils/erasers.
 - Structure of test will be described today.

Compactness

Definition (Heine-Borel Property)

A set $E \subseteq \mathbb{R}$ is said to have the **Heine-Borel property** if every open cover of E can be reduced to a finite subcover. That is, if \mathcal{U} is an open cover of E , then there exists a finite subfamily $\{U_1, U_2, \dots, U_n\} \subseteq \mathcal{U}$, such that $E \subseteq U_1 \cup U_2 \cup \dots \cup U_n$.

When does any open cover of a set E have a finite subcover?

Theorem (Heine-Borel Theorem)

A set $E \subseteq \mathbb{R}$ has the *Heine-Borel property* iff E is both *closed* and *bounded*.

(Textbook (TBB) pp. 249–250)

Compactness

Definition (Compact Set)

A set $E \subseteq \mathbb{R}$ is said to be **compact** if it has any of the following equivalent properties:

- 1 E is **closed** and bounded.
- 2 E has the **Bolzano-Weierstrass property**.
- 3 E has the **Heine-Borel property**.

Note: In spaces other than \mathbb{R} , these three properties are not necessarily equivalent. Usually the **Heine-Borel property** is taken as the definition of compactness.

Compactness

Example

Prove that the interval $(0, 1]$ is not compact by showing that it is not closed or not bounded.

(solution on board)

Example

Prove that the interval $(0, 1]$ is not compact by showing that it does not have the **Bolzano-Weierstrass property**.

(solution on board)

Example

Prove that the interval $(0, 1]$ is not compact by showing that it does not have the **Heine-Borel property**.

(solution on board)

Compactness

Example (Classic non-trivial compactness argument)

Let E be a **compact** subset of \mathbb{R} . Prove that if $f : E \rightarrow \mathbb{R}$ is **locally bounded** on E then f is **bounded** on E .

(solution on board)

Bolzano-Weierstrass approach: Textbook (TBB) p. 242

Heine-Borel approach: Textbook (TBB) p. 251

Example (Converse of above example)

Let $E \subseteq \mathbb{R}$. If every function $f : E \rightarrow \mathbb{R}$ that is **locally bounded** on E is **bounded** on E , then E is **compact**.

(solution on board)

Note: Contrapositive of converse is: If $E \subseteq \mathbb{R}$ is not **compact** then $\exists f : E \rightarrow \mathbb{R}$ \nrightarrow f is **locally bounded** on E but not **bounded** on E . •

Complements and Closures problem

Example

How many distinct sets can be obtained from $E = [0, 1]$ by applying the complement and closure operations?

Consider this sequence of sets: $E_1 = [0, 1]$,
 $E_2 = E_1^c = (-\infty, 0) \cup (1, \infty)$, $E_3 = \overline{E_2} = (-\infty, 0] \cup [1, \infty)$,
 $E_4 = E_3^c = (0, 1)$, $E_5 = \overline{E_4} = E_1$.

Does this prove the answer is 4?

Extra Challenge Problem

If $E \subseteq \mathbb{R}$, how many distinct sets can be obtained by taking complements or closures of E and its successors? Put another way, if $\{E_n\}$ is a sequence of sets produced by taking the complement or closure of the previous set, how many distinct sets can such a sequence contain? If the answer is finite, find a set E that generates the maximum number in this way.

Test #1

What you need to know:

- Everything discussed in class, including all definitions/concepts and theorems/lemmas/corollaries.
- Everything in assignments and solutions to assignments.
Make sure you fully understand all the solutions to all the problems in all the assignments.
- Most—but not all—of the material that you are responsible for is covered in chapters 1, 2 and 4 of the textbook. You are not responsible for material in the textbook that was not covered in lectures or assignments.
- It is essential that you understand how to use the definitions and theorems to construct proofs.

Test #1

Other comments:

- You will not be asked to list all the axioms for the the real number system. BUT, you must be able to state the *completeness axiom* concerning least upper bounds.
- You should do last year's Test #1 for practice.
- There are many additional problems that would be good to try for practice in chapters 1, 2 and 4 of the textbook.
- Structure of the test:
 - Question 1 is the same as last year.
 - Questions 2, 3 and 4 are different types of multiple choice questions, e.g., "TRUE" or "FALSE" or "Always TRUE", "Sometimes TRUE" or "Never TRUE".
 - Questions 5 and 6 and tables to fill in.
 - Questions 7, 8 and 9 require proofs.

Student Name: _____

Student Number: _____

Mathematics 3A03 — Real Analysis I

TERM TEST #1 — 23 October 2017

Duration: 90 minutes

Notes:

- No calculators, notes, scrap paper, or aids of any kind are permitted.
- This test consists of **8 pages** (i.e., **4 double-sided pages**). There are **9 questions** in total. Bring any discrepancy to the attention of your instructor or invigilator.
- All questions are to be answered on this test paper. The final page is blank to provide extra space if needed.
- The first 6 questions do not require any justification for your answers. For these, you will be assessed on your answers only. *Do not justify your answers to these questions.*
- Always write clearly. An answer that cannot be deciphered cannot be marked.
- The marking scheme is indicated in the margin. The maximum total mark is 50.

Question	Mark
1	
2	
3	
4	
5	
Subtotal	

Question	Mark
6	
7	
8	
9	
Subtotal	

Total	
-------	--

Student Name: _____

Student Number: _____

Mathematics 3A03 — Real Analysis I

TERM TEST #1 — 23 October 2017

Duration: 90 minutes

Notes:

- No calculators, notes, scrap paper, or aids of any kind are permitted.
- This test consists of **8 pages** (*i.e.*, **4 double-sided pages**). There are **9 questions** in total. Bring any discrepancy to the attention of your instructor or invigilator.
- All questions are to be answered on this test paper. The final page is blank to provide extra space if needed.
- The first 6 questions do not require any justification for your answers. For these, you will be assessed on your answers only. *Do not justify your answers to these questions.*
- Always write clearly. An answer that cannot be deciphered cannot be marked.
- The marking scheme is indicated in the margin. The maximum total mark is 50.

Question	Mark

Question	Mark