

**25** Differentiation

**26** Differentiation II

# Differentiation



Mathematics  
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

# Mathematics 3A03 Real Analysis I


Instructor: David Earn

Lecture 25  
Differentiation  
Wednesday 8 November 2017

# Announcements

- **Assignment 5** has been posted on the course wiki. The assignment is due on Monday 20 Nov 2017 at 2:25pm (remember cover sheet!)

# The Derivative



# The Derivative

## Definition (Derivative)

Let  $f$  be defined on an interval  $I$  and let  $x_0 \in I$ . The **derivative** of  $f$  at  $x_0$ , denoted by  $f'(x_0)$ , is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If  $f'(x_0)$  is finite we say that  $f$  is **differentiable** at  $x_0$ . If  $f$  is differentiable at every point of a set  $E \subseteq I$ , we say that  $f$  is differentiable on  $E$ . If  $E$  is all of  $I$ , we simply say that  $f$  is a **differentiable function**.

Note: “Differentiable” and “a derivative exists” always mean that the derivative is finite.

# The Derivative

## Example

$f(x) = x^2$ . Find  $f'(2)$ .

$$f'(2) = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \lim_{x \rightarrow 2} \frac{(x + 2)(x - 2)}{x - 2} = \lim_{x \rightarrow 2} x + 2 = 4$$


### Note:

- In the first two limits, we must have  $x \neq 2$ .
- But in the third limit, we just plug in  $x = 2$ .
- Two things are equal, but in one  $x \neq 2$  and in the other  $x = 2$ .
- Good illustration of why it is important to define the meaning of limits rigorously.

# The Derivative

## Example

Let  $f$  be defined in a neighbourhood  $I$  of 0, and suppose  $|f(x)| \leq x^2$  for all  $x \in I$ . Is  $f$  necessarily differentiable at 0? e.g.,



(solution on board)



# The Derivative

## Definition (One-sided derivatives)

Let  $f$  be defined on an interval  $I$  and let  $x_0 \in I$ . The **right-hand derivative** of  $f$  at  $x_0$ , denoted by  $f'_+(x_0)$ , is the limit

$$f'_+(x_0) = \lim_{x \rightarrow x_0^+} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this one-sided limit exists or is infinite.

Similarly, the **left-hand derivative** of  $f$  at  $x_0$ , denoted by  $f'_-(x_0)$ , is the limit

$$f'_-(x_0) = \lim_{x \rightarrow x_0^-} \frac{f(x) - f(x_0)}{x - x_0}.$$


### Note:

If  $x_0 \in I^\circ$  then  $f$  is differentiable at  $x_0$  iff  $f'_+(x_0) = f'_-(x_0) \neq \pm\infty$ .

# The Derivative

## Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$



- Same slope from left and right. Why isn't  $f$  differentiable???
- $\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^+} f'(x) = \lim_{x \rightarrow 0} f'(x) = 1.$
- $f'_-(0) = f'_+(0) = f'(0) = \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \infty.$

# The Derivative

- Higher derivatives: we write
  - $f'' = (f')'$  if  $f'$  is differentiable;
  - $f^{(n+1)} = (f^{(n)})'$  if  $f^{(n)}$  is differentiable.
- Other standard notation for derivatives:

$$\frac{df}{dx} = f'(x)$$

$$D = \frac{d}{dx}$$

$$D^n f(x) = \frac{d^n f}{dx^n} = f^{(n)}(x)$$

# The Derivative

## Theorem (Differentiable $\implies$ continuous)

If  $f$  is defined in a neighbourhood  $I$  of  $x_0$  and  $f$  is differentiable at  $x_0$  then  $f$  is continuous at  $x_0$ .

### Proof.

Must show  $\lim_{x \rightarrow x_0} f(x) = f(x_0)$ , i.e.,  $\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = 0$ .

$$\begin{aligned}\lim_{x \rightarrow x_0} (f(x) - f(x_0)) &= \lim_{x \rightarrow x_0} \left( \frac{f(x) - f(x_0)}{x - x_0} \times (x - x_0) \right) \\ &= \lim_{x \rightarrow x_0} \left( \frac{f(x) - f(x_0)}{x - x_0} \right) \times \lim_{x \rightarrow x_0} (x - x_0) \\ &= f'(x_0) \times 0 = 0,\end{aligned}$$

where we have used the theorem on the algebra of limits.  $\square$

# The Derivative

## Theorem (Algebra of derivatives)

*Suppose  $f$  and  $g$  are defined on an interval  $I$  and  $x_0 \in I$ . If  $f$  and  $g$  are differentiable at  $x_0$  then  $f + g$  and  $fg$  are differentiable at  $x_0$ . If, in addition,  $g(x_0) \neq 0$  then  $f/g$  is differentiable at  $x_0$ . Under these conditions:*

**1**  $(cf)'(x_0) = cf'(x_0)$  for all  $c \in \mathbb{R}$ ;

**2**  $(f + g)'(x_0) = (f' + g')(x_0)$ ;

**3**  $(fg)'(x_0) = (f'g + fg')(x_0)$ ;

**4**  $\left(\frac{f}{g}\right)'(x_0) = \left(\frac{gf' - fg'}{g^2}\right)(x_0) \quad (g(x_0) \neq 0).$

(Textbook (TBB) [Theorem 7.7, p. 408](#))

# The Derivative

## Theorem (Chain rule)

*Suppose  $f$  is defined in a neighbourhood  $U$  of  $x_0$  and  $g$  is defined in a neighbourhood  $V$  of  $f(x_0)$  such that  $f(U) \subseteq V$ . If  $f$  is differentiable at  $x_0$  and  $g$  is differentiable at  $f(x_0)$  then the composite function  $h = g \circ f$  is differentiable at  $x_0$  and*

$$h'(x_0) = (g \circ f)'(x_0) = g'(f(x_0))f'(x_0).$$

(Textbook (TBB) §7.3.2)

TBB provide a very good motivating discussion of this proof, which is quite technical.


# The Derivative

## Theorem (Derivative at local extrema)

Let  $f : (a, b) \rightarrow \mathbb{R}$ . If  $x$  is a maximum or minimum point of  $f$  in  $(a, b)$ , and  $f$  is differentiable at  $x$ , then  $f'(x) = 0$ .

(Textbook (TBB) [Theorem 7.18](#), p. 424)

Note:  $f$  need not be differentiable or even continuous at other points.





Mathematics  
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

# Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 26  
Differentiation II  
Friday 10 November 2017



# Announcements

- [Assignment 5](#) has been posted on the course wiki. The assignment is due on Monday 20 Nov 2017 at 2:25pm (remember cover sheet!)

# Survey results

Table 14: The status quo is that there are two lecture sections in which the two instructors deliver the same material simultaneously, and four tutorial sections that are designed and led by the two TAs (who present problems at the blackboard). Which of the following options would you prefer for the remainder of the course?

P01	Preference	n	percent
1	Maintain the status quo	46	57.50
2	Maintain the status quo for lectures, but run tutorials in a different style where students work on problems in groups and the TA circulates giving advice on constructing proofs	15	18.80
3	Have everyone attend the same lecture section (with the two instructors alternating topics) but instructors and TAs co-design and co-run tutorials in a more "group work" fashion	7	8.75
4	NA's	12	15.00

# Last time...

- Definition of the derivative.
- Proved differentiable  $\implies$  continuous.
- Discussed algebra of derivatives and chain rule.
- Proved derivative is zero at extrema.
- Defined one-sided derivatives
  - Example

# The Mean Value Theorem

## Theorem (Rolle's theorem)

*If  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , and  $f(a) = f(b)$ , then there exists  $x \in (a, b)$  such that  $f'(x) = 0$ .*

(solution on board)

(Textbook (TBB) [Theorem 7.19](#), p. 427)


## Theorem (Mean value theorem)

*If  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$  then there exists  $x \in (a, b)$  such that*

$$f'(x) = \frac{f(b) - f(a)}{b - a}.$$

# The Mean Value Theorem

*Idea for proof:*



**Proof.**

Apply [Rolle's theorem](#) to

$$h(x) = f(x) - \left[ f(a) + \left( \frac{f(b) - f(a)}{b - a} \right) (x - a) \right].$$

□

# The Mean Value Theorem

## Example

$f'(x) > 0$  on an interval  $I \implies f$  strictly increasing on  $I$ .

*Proof:*

Suppose  $x_1, x_2 \in I$  and  $x_1 < x_2$ . We must show  $f(x_1) < f(x_2)$ .

Since  $f'(x)$  exists for all  $x \in I$ ,  $f$  is certainly differentiable on the closed subinterval  $[x_1, x_2]$ .

Hence by the **Mean Value Theorem**  $\exists x_* \in (x_1, x_2)$  such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(x_*).$$

But  $x_2 - x_1 > 0$  and since  $x_* \in I$ , we know  $f'(x_*) > 0$ .

$\therefore f(x_2) - f(x_1) > 0$ , *i.e.*,  $f(x_1) < f(x_2)$ . □

# Intermediate value property for derivatives

## Theorem (Darboux's Theorem: IVP for derivatives)

If  $f$  is differentiable on an interval  $I$  then its derivative  $f'$  has the *intermediate value property* on  $I$ .

### Notes:

- *Equivalent (contrapositive) statement of Darboux's theorem:*  
If a function does not have the *intermediate value property* on  $I$  then it is impossible that it is the derivative of any function on  $I$ .
- It is  $f'$ , not  $f$ , that is claimed to have the *intermediate value property* in Darboux's theorem. This theorem does not follow from the standard *intermediate value theorem* because the derivative  $f'$  is not necessarily continuous.
- Darboux's theorem implies that a derivative cannot have jump or removable discontinuities. Any discontinuity of a derivative must be *essential*. Recall example of a *discontinuous function with IVP*.

# Intermediate value property for derivatives

## Proof of Darboux's Theorem.

Consider  $a, b \in I$  with  $a < b$ .

Suppose first that  $f'(a) < 0 < f'(b)$ . We will show  $\exists x \in (a, b)$  such that  $f'(x) = 0$ . Since  $f'$  exists on  $[a, b]$ , we must have  $f$  continuous on  $[a, b]$ , so the **Extreme Value Theorem** implies that  $f$  attains its minimum at some point  $x \in [a, b]$ . This minimum point cannot be an endpoint of  $[a, b]$  ( $x \neq a$  because  $f'(a) < 0$  and  $x \neq b$  because  $f'(b) > 0$ ).

Therefore,  $x \in (a, b)$ . But  $f$  is differentiable everywhere in  $(a, b)$ , so, by the **theorem on the derivative at local extrema**, we must have  $f'(x) = 0$ .

Now suppose more generally that  $f'(a) < K < f'(b)$ . Let  $g(x) = f(x) - Kx$ . Then  $g$  is differentiable on  $I$  and  $g'(x) = f'(x) - K$  for all  $x \in I$ . In addition,  $g'(a) = f'(a) - K < 0$  and  $g'(b) = f'(b) - K > 0$ , so by the argument above,  $\exists x \in (a, b)$  such that  $g'(x) = 0$ , i.e.,  $f'(x) - K = 0$ , i.e.,  $f'(x) = K$ .

The case  $f'(a) > K > f'(b)$  is similar. □



# Intermediate value property for derivatives

Example  $(f'(x) \neq 0 \forall x \implies f \nearrow \text{ or } \searrow)$

If  $f$  is differentiable on an interval  $I$  and  $f'(x) \neq 0$  for all  $x \in I$  then  $f$  is either increasing or decreasing on the entire interval  $I$ .

*Proof:*

Suppose  $\exists a, b \in I$  such that  $f'(a) < 0$  and  $f'(b) > 0$ .

Then, from [Darboux's theorem](#),  $\exists c \in I$  such that  $f'(c) = 0$ .  $\implies \Leftarrow$

$\therefore$  Either " $\exists a \in I \wedge f'(a) < 0$ " is FALSE  
 or " $\exists b \in I \wedge f'(b) > 0$ " is FALSE.

$\therefore$  Since we know  $f'(x) \neq 0 \forall x \in I$ , it must be that  
either  $f'(x) > 0 \forall x \in I$  or  $f'(x) < 0 \forall x \in I$ ,  
*i.e.*, either  $f$  is increasing on  $I$  or decreasing on  $I$ . □

# The Derivative of an Inverse

## Theorem (Inverse function theorem)

If  $f$  is differentiable on an interval  $I$  and  $f'(x) \neq 0 \forall x \in I$ , then

1  $f$  is one-to-one on  $I$ ;

2  $f^{-1}$  is differentiable on  $J = f(I)$ ;

3  $(f^{-1})'(f(x)) = \frac{1}{f'(x)}$  for all  $x \in I$ ,

i.e.,  $(f^{-1})'(y) = \frac{1}{f'((f^{-1}(y)))}$  for all  $y \in J$ .

(Textbook (TBB) [Theorem 7.32](#), p. 445)

# The Derivative of an Inverse

Key insights for proof of inverse function theorem:

- Darboux's theorem  $\implies f$  is 1 : 1 on  $I$

- If  $y = f(x)$  and  $y_0 = f(x_0)$

then  $x = f^{-1}(y)$  and  $x_0 = f^{-1}(y_0)$ ,

$$\begin{aligned} \text{so } \frac{f^{-1}(y) - f^{-1}(y_0)}{y - y_0} &= \frac{x - x_0}{f(x) - f(x_0)} \\ &= \frac{1}{\frac{f(x) - f(x_0)}{x - x_0}}. \end{aligned}$$

- Since  $f$  continuous at  $x_0$ , we know  $x \rightarrow x_0 \implies y \rightarrow y_0$ .
- But we need  $y \rightarrow y_0 \implies x \rightarrow x_0$ , i.e.,  $f^{-1}$  continuous at  $y_0$ .
- In fact,  $f$  continuous and either  $\nearrow$  or  $\searrow$  on  $I \implies f^{-1}$  continuous on  $J = f(I)$ . (more generally, cf. [Invariance of Domain](#) thm)