

26 Integration

27 Integration II

28 Integration III



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$


Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 26
Integration
Friday 8 November 2019

Integration


Integration



- “Area of region $R(f, a, b)$ ” is actually a very subtle concept.
- We will only scratch the surface of it.
- Textbook presentation of integral is different (but equivalent).


Our treatment is closer to that in M. Spivak “Calculus” (2008).

Integration




- Contribution to “area of $R(f, a, b)$ ” is positive or negative depending on whether f is positive or negative.


Lower sum




Upper sum




Lower and upper sums




Lower and upper sums




Lower and upper sums




Lower and upper sums



Lower and upper sums



Lower and upper sums



Rigorous development of the integral

Definition (Partition)

Let $a < b$. A **partition** of the interval $[a, b]$ is a finite collection of points in $[a, b]$, one of which is a , and one of which is b .

We normally label the points in a partition

$$a = t_0 < t_1 < \cdots < t_{n-1} < t_n = b,$$

so the i th subinterval in the partition is

$$[t_{i-1}, t_i].$$

Rigorous development of the integral

Definition (Lower and upper sums)

Suppose f is bounded on $[a, b]$ and $P = \{t_0, \dots, t_n\}$ is a **partition** of $[a, b]$. Let

$$m_i = \inf \{ f(x) : x \in [t_{i-1}, t_i] \},$$

$$M_i = \sup \{ f(x) : x \in [t_{i-1}, t_i] \}.$$

The **lower sum** of f for P , denoted by $L(f, P)$, is defined as

$$L(f, P) = \sum_{i=1}^n m_i(t_i - t_{i-1}).$$

The **upper sum** of f for P , denoted by $U(f, P)$, is defined as

$$U(f, P) = \sum_{i=1}^n M_i(t_i - t_{i-1}).$$

Rigorous development of the integral

Relationship between motivating sketch and rigorous definition of lower and upper sums:

- The **lower and upper sums** correspond to the total areas of rectangles lying below and above the graph of f in our **motivating sketch**.
- However, these sums have been defined precisely without any appeal to a concept of “area”.
- The requirement that f be bounded on $[a, b]$ is essential in order that all the m_i and M_i be well-defined.
- It is also essential that the m_i and M_i be defined as inf's and sup's (rather than maxima and minima) because f was not assumed continuous.

Rigorous development of the integral

Relationship between motivating sketch and rigorous definition of lower and upper sums:

- Since $m_i \leq M_i$ for each i , we have

$$m_i(t_i - t_{i-1}) \leq M_i(t_i - t_{i-1}). \quad i = 1, \dots, n.$$

∴ For any partition P of $[a, b]$ we have

$$L(f, P) \leq U(f, P),$$

because

$$L(f, P) = \sum_{i=1}^n m_i(t_i - t_{i-1}),$$
$$U(f, P) = \sum_{i=1}^n M_i(t_i - t_{i-1}).$$

Poll

- Go to https://www.childsmath.ca/childsa/forms/main_login.php
- Click on [Math 3A03](#)
- Click on [Take Class Poll](#)
- Fill in poll **Lecture 26: Lower and Upper Sums**
- .

Rigorous development of the integral

Relationship between motivating sketch and rigorous definition of lower and upper sums:

- More generally, if P_1 and P_2 are any two partitions of $[a, b]$, it ought to be true that

$$L(f, P_1) \leq U(f, P_2),$$

because $L(f, P_1)$ should be \leq area of $R(f, a, b)$, and $U(f, P_2)$ should be \geq area of $R(f, a, b)$.

- But “ought to” and “should be” prove nothing, especially since we haven’t yet even defined “area of $R(f, a, b)$ ”.
- Before we can *define* “area of $R(f, a, b)$ ”, we need to prove that $L(f, P_1) \leq U(f, P_2)$ for any partitions $P_1, P_2 \dots$



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 27
Integration II
Tuesday 12 November 2019

Poll

- Go to https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Lecture 27: Assignment 3 Awareness**
- .


Announcements

- **Assignment 4** was due before class today.
- **Assignment 5** is due on **Thursday 21 November 2019 @ 2:25pm** via **crowdmark**.
- **Math 3A03 Test #2**
Tuesday 26 November 2019, 5:30–7:00pm, in JHE 264
- **Assignment 6** will be due on **Tuesday 3 December 2019 @ 2:25pm** via **crowdmark**.
- **Math 3A03 Final Exam:** Fri 6 Dec 2019, 9:00am–11:30am
Location: MDCL 1105

Rigorous development of the integral

Lemma

If *partition* $P \subseteq$ *partition* Q (i.e., if every point of P is also in Q), then $L(f, P) \leq L(f, Q)$ and $U(f, P) \geq U(f, Q)$.



Rigorous development of the integral

Proof of Lemma

As a first step, consider the special case in which the finer partition Q contains only one more point than P :

$$P = \{t_0, \dots, t_n\},$$

$$Q = \{t_0, \dots, t_{k-1}, u, t_k, \dots, t_n\},$$

where

$$a = t_0 < t_1 < \dots < t_{k-1} < u < t_k < \dots < t_n = b.$$

Let

$$m' = \inf \{ f(x) : x \in [t_{k-1}, u] \},$$

$$m'' = \inf \{ f(x) : x \in [u, t_k] \}.$$

... continued ...

Rigorous development of the integral

Proof of Lemma (cont.)

Then
$$L(f, P) = \sum_{i=1}^n m_i(t_i - t_{i-1}),$$

and
$$L(f, Q) = \sum_{i=1}^{k-1} m_i(t_i - t_{i-1}) + m'(u - t_{k-1}) \\ + m''(t_k - u) + \sum_{i=k+1}^n m_i(t_i - t_{i-1}).$$

\therefore To prove $L(f, P) \leq L(f, Q)$, it is enough to show

$$m_k(t_k - t_{k-1}) \leq m'(u - t_{k-1}) + m''(t_k - u).$$

... continued ...

Rigorous development of the integral

Proof of Lemma (cont.)

Now note that since

$$\{f(x) : x \in [t_{k-1}, u]\} \subseteq \{f(x) : x \in [t_{k-1}, t_k]\},$$

the RHS might contain some additional *smaller* numbers, so we must have

$$\begin{aligned} m_k &= \inf \{f(x) : x \in [t_{k-1}, t_k]\} \\ &\leq \inf \{f(x) : x \in [t_{k-1}, u]\} = m'. \end{aligned}$$

Thus, $m_k \leq m'$, and, similarly, $m_k \leq m''$.

$$\begin{aligned} \therefore m_k(t_k - t_{k-1}) &= m_k(t_k - u + u - t_{k-1}) \\ &= m_k(u - t_{k-1}) + m_k(t_k - u) \\ &\leq m'(u - t_{k-1}) + m''(t_k - u), \end{aligned}$$

... continued ...

Rigorous development of the integral

Proof of Lemma (cont.)

which proves (in this special case where Q contains only one more point than P) that $L(f, P) \leq L(f, Q)$.

We can now prove the general case by adding one point at a time.

If Q contains ℓ more points than P , define a sequence of partitions

$$P = P_0 \subset P_1 \subset \cdots \subset P_\ell = Q$$

such that P_{j+1} contains exactly one more point than P_j . Then

$$L(f, P) = L(f, P_0) \leq L(f, P_1) \leq \cdots \leq L(f, P_\ell) = L(f, Q),$$

so $L(f, P) \leq L(f, Q)$.

(Proving $U(f, P) \geq U(f, Q)$ is similar: check!)



Rigorous development of the integral

Theorem (Partition Theorem)

Let P_1 and P_2 be any two partitions of $[a, b]$. If f is bounded on $[a, b]$ then

$$L(f, P_1) \leq U(f, P_2).$$

Proof.

This is a straightforward consequence of the [partition lemma](#).

Let $P = P_1 \cup P_2$, i.e., the partition obtained by combining all the points of P_1 and P_2 .

Then

$$L(f, P_1) \leq L(f, P) \leq U(f, P) \leq U(f, P_2).$$



Rigorous development of the integral

Important inferences that follow from the [partition theorem](#):

- For any partition P' , the upper sum $U(f, P')$ is an upper bound for the set of all lower sums $L(f, P)$.

$$\therefore \sup \{L(f, P) : P \text{ a partition of } [a, b]\} \leq U(f, P') \quad \forall P'$$

$$\therefore \sup \{L(f, P)\} \leq \inf \{U(f, P)\}$$

\therefore For any partition P' ,

$$L(f, P') \leq \sup \{L(f, P)\} \leq \inf \{U(f, P)\} \leq U(f, P')$$

- If $\sup \{L(f, P)\} = \inf \{U(f, P)\}$ then we can define “area of $R(f, a, b)$ ” to be this number.

- Is it possible that $\sup \{L(f, P)\} < \inf \{U(f, P)\}$?

Rigorous development of the integral

Example

$\exists? f : [a, b] \rightarrow \mathbb{R}$ such that $\sup \{L(f, P)\} < \inf \{U(f, P)\}$

Let

$$f(x) = \begin{cases} 1 & x \in \mathbb{Q} \cap [a, b], \\ 0 & x \in \mathbb{Q}^c \cap [a, b]. \end{cases}$$

If $P = \{t_0, \dots, t_n\}$ then $m_i = 0$ ($\because [t_{i-1}, t_i] \cap \mathbb{Q}^c \neq \emptyset$),
and $M_i = 1$ ($\because [t_{i-1}, t_i] \cap \mathbb{Q} \neq \emptyset$).

$\therefore L(f, P) = 0$ and $U(f, P) = b - a$ for any partition P .

$\therefore \sup \{L(f, P)\} = 0 < b - a = \inf \{U(f, P)\}$. □

Can we define “area of $R(f, a, b)$ ” for such a weird function?

Yes, but not in this course!

Rigorous development of the integral

Definition (Integrable)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *integrable* on $[a, b]$ if it is bounded on $[a, b]$ and

$$\begin{aligned} & \sup\{L(f, P) : P \text{ a partition of } [a, b]\} \\ & = \inf\{U(f, P) : P \text{ a partition of } [a, b]\}. \end{aligned}$$

In this case, this common number is called the *integral* of f on $[a, b]$ and is denoted

$$\int_a^b f$$

Note: If f is integrable then for any partition P we have

$$L(f, P) \leq \int_a^b f \leq U(f, P),$$

and $\int_a^b f$ is the unique number with this property.

Rigorous development of the integral

- *Notation:*

$$\int_a^b f(x) dx \quad \text{means precisely the same as} \quad \int_a^b f.$$

- The symbol “ dx ” has no meaning in isolation just as “ $x \rightarrow$ ” has no meaning except in $\lim_{x \rightarrow a} f(x)$.
- It is not clear from the definition which functions are **integrable**.
- The definition of the **integral** does not itself indicate how to compute the integral of any given **integrable** function. So far, without a lot more effort we can't say much more than these two things:
 - 1 If $f(x) \equiv c$ then f is **integrable** on $[a, b]$ and $\int_a^b f = c \cdot (b - a)$.
 - 2 The **weird example** function is not **integrable**.

Rigorous development of the integral

- A function that is **integrable** according to our definition is usually said to be **Riemann integrable**, to distinguish this definition from other definitions of integrability.
- In Math 4A03 you will define “Lebesgue integrable”, a more subtle concept that makes it possible to attach meaning to “area of $R(f, a, b)$ ” for the **weird example** function (among others), and to precisely characterize functions that are Riemann integrable.

Rigorous development of the integral

Theorem (Equivalent condition for integrability)

A *bounded* function $f : [a, b] \rightarrow \mathbb{R}$ is *integrable* on $[a, b]$ iff for all $\varepsilon > 0$ there is a partition P of $[a, b]$ such that

$$U(f, P) - L(f, P) < \varepsilon.$$

Proof.

2016 Assignment 5. □

Note: This theorem is just a restatement of the definition of integrability. It is often more convenient to work with $\varepsilon > 0$ than with sup's and inf's.



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 28
Integration III
Thursday 14 November 2019

Last time...

Rigorous development of integral:

- Definition: **integrable**.
- Example: **non-integrable function**.
- Theorem: Equivalent " ϵ - P " definition of integrable.

Poll

- Go to https://www.childsmath.ca/childsaf/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Lecture 28: Integrable vs Continuous vs Differentiable**
- .

Integral theorems

Theorem

If f is continuous on $[a, b]$ then f is *integrable* on $[a, b]$.

Rough work to prepare for proof:

$$U(f, P) - L(f, P) = \sum_{i=1}^n (M_i - m_i)(t_i - t_{i-1})$$

Given $\varepsilon > 0$, choose a partition P that is so fine that $M_i - m_i < \varepsilon$ for all i . Then

$$U(f, P) - L(f, P) < \varepsilon \sum_{i=1}^n (t_i - t_{i-1}) = \varepsilon(b - a).$$

Not quite what we want. So choose the partition P such that $M_i - m_i < \varepsilon/(b - a)$ for all i . To get that, choose P such that

$$|f(x) - f(y)| < \frac{\varepsilon}{2(b - a)} \quad \text{if } |x - y| < \max_{1 \leq i \leq n} (t_i - t_{i-1}),$$

which we can do because f is uniformly continuous on $[a, b]$.

Integral theorems

Proof that continuous \implies integrable

Since f is continuous on the compact set $[a, b]$, it is bounded on $[a, b]$ (which is the first requirement to be **integrable** on $[a, b]$).

Also, since f is continuous on the compact set $[a, b]$, it is uniformly continuous on $[a, b]$. $\therefore \forall \varepsilon > 0 \exists \delta > 0$ such that $\forall x, y \in [a, b]$,

$$|x - y| < \delta \implies |f(x) - f(y)| < \frac{\varepsilon}{2(b - a)}.$$

Now choose a partition of $[a, b]$ such that the length of each subinterval $[t_{i-1}, t_i]$ is less than δ , i.e., $t_i - t_{i-1} < \delta$. Then, for any $x, y \in [t_{i-1}, t_i]$ we have $|x - y| < \delta$ and therefore

... continued ...

Integral theorems

Proof that continuous \implies integrable (cont.)

$$|f(x) - f(y)| < \frac{\varepsilon}{2(b-a)} \quad \forall x, y \in [t_{i-1}, t_i].$$

$$\therefore M_i - m_i \leq \frac{\varepsilon}{2(b-a)} < \frac{\varepsilon}{b-a} \quad i = 1, \dots, n.$$

Since this is true for all i , it follows that

$$\begin{aligned} U(f, P) - L(f, P) &= \sum_{i=1}^n (M_i - m_i)(t_i - t_{i-1}) \\ &< \frac{\varepsilon}{b-a} \sum_{i=1}^n (t_i - t_{i-1}) = \frac{\varepsilon}{b-a} (b-a) = \varepsilon. \end{aligned}$$

□

Properties of the integral

Theorem (Integral segmentation)

Let $a < c < b$. If f is *integrable* on $[a, b]$, then f is *integrable* on $[a, c]$ and on $[c, b]$. Conversely, if f is *integrable* on $[a, c]$ and $[c, b]$ then f is *integrable* on $[a, b]$. Finally, if f is *integrable* on $[a, b]$ then

$$\int_a^b f = \int_a^c f + \int_c^b f. \quad (\heartsuit)$$

(a good exercise)

This theorem motivates these definitions:

$$\int_a^a f = 0 \quad \text{and} \quad \int_a^b f = -\int_b^a f \quad \text{if } a > b.$$

Then (\heartsuit) holds for any $a, b, c \in \mathbb{R}$.

Properties of the integral

Theorem (Algebra of integrals – a.k.a. \int_a^b is a linear operator)

If f and g are *integrable* on $[a, b]$ and $c \in \mathbb{R}$ then $f + g$ and cf are *integrable* on $[a, b]$ and

$$\mathbf{1} \quad \int_a^b (f + g) = \int_a^b f + \int_a^b g;$$

$$\mathbf{2} \quad \int_a^b cf = c \int_a^b f.$$

(proofs are relatively easy; good exercises)

Theorem (Integral of a product)

If f and g are *integrable* on $[a, b]$ then fg is *integrable* on $[a, b]$.

(proof is much harder; tough exercise)

Properties of the integral

Lemma (Integral bounds)

Suppose f is integrable on $[a, b]$. If $m \leq f(x) \leq M$ for all $x \in [a, b]$ then

$$m(b - a) \leq \int_a^b f \leq M(b - a).$$

Proof.

For any partition P , we must have $m \leq m_i \forall i$ and $M \geq M_i \forall i$.

$$\therefore m(b - a) \leq L(f, P) \leq U(f, P) \leq M(b - a) \quad \forall P$$

$$\therefore m(b - a) \leq \sup\{L(f, P)\} = \int_a^b f = \inf\{U(f, P)\} \leq M(b - a).$$



Properties of the integral

Theorem (Integrals are continuous)

If f is *integrable* on $[a, b]$ and F is defined on $[a, b]$ by

$$F(x) = \int_a^x f,$$

then F is continuous on $[a, b]$.

Proof

Let's first consider $x_0 \in [a, b)$ and show F is continuous from above at x_0 , i.e., $\lim_{x \rightarrow x_0^+} F(x) = F(x_0)$. If $x \in (x_0, b]$ then

$$(\heartsuit) \quad \implies \quad F(x) - F(x_0) = \int_a^x f - \int_a^{x_0} f = \int_{x_0}^x f. \quad (*)$$

... continued ...

Properties of the integral

Proof (cont.)

Since f is **integrable** on $[a, b]$, it is bounded on $[a, b]$, so $\exists M > 0$ such that

$$-M \leq f(x) \leq M \quad \forall x \in [a, b],$$

from which the **integral bounds lemma** implies

$$-M(x - x_0) \leq \int_{x_0}^x f \leq M(x - x_0),$$

$$\therefore (*) \implies -M(x - x_0) \leq F(x) - F(x_0) \leq M(x - x_0).$$

\therefore For any $\varepsilon > 0$ we can ensure $|F(x) - F(x_0)| < \varepsilon$ by requiring $0 \leq x - x_0 < \varepsilon/M$, which proves $\lim_{x \rightarrow x_0^+} F(x) = F(x_0)$.

A similar argument starting from $x_0 \in (a, b]$ and $x \in [a, x_0)$ yields $\lim_{x \rightarrow x_0^-} F(x) = F(x_0)$. Thus, “integrals are continuous”. \square