

13 Topology of \mathbb{R} I

14 Topology of \mathbb{R} II

15 Topology of \mathbb{R} III



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 13
Topology of \mathbb{R}^1
Tuesday 1 October 2019

THINKING ABOUT GRADUATE SCHOOL?

JOIN US TO FIND OUT MORE AT THE GRAD
INFO SESSION!

WHEN: THURSDAY OCTOBER 3, 2019

TIME: 5:30PM – 7:00PM

WHERE: HH/305 AND THE MATH CAFÉ

Matheus Grasselli will give general advice on applying to grad school.

Shui Feng will talk about graduate programs particular to statistics.

Tom Hurd will talk about graduate opportunities in financial math including PhiMac.

Miroslav Lovric will give tips about applying to teachers' college.

PIZZA will be served! See you there!



Announcements

- **Assignment 3** is posted, but more problems will be added in a few days. **Due Tuesday 22 October 2019 at 2:25pm via crowdmark.**

Topology of \mathbb{R}

Intervals



Open interval:

$$(a, b) = \{x : a < x < b\}$$


Closed interval:

$$[c, d] = \{x : c \leq x \leq d\}$$

Half-open interval:

$$(e, f] = \{x : e < x \leq f\}$$

Interior point




Definition (Interior point)

If $E \subseteq \mathbb{R}$ then x is an *interior point* of E if x lies in an open interval that is contained in E , i.e., $\exists c > 0$ such that $(x - c, x + c) \subset E$.

Interior point examples

Set E	Interior points?
$(-1, 1)$	Every point
$[0, 1]$	Every point <i>except the endpoints</i>
\mathbb{N}	\nexists
\mathbb{R}	Every point
\mathbb{Q}	\nexists
$(-1, 1) \cup [0, 1]$	Every point <i>except 1</i>
$(-1, 1) \setminus \{\frac{1}{2}\}$	Every point


Neighbourhood



Definition (Neighbourhood)

A *neighbourhood* of a point $x \in \mathbb{R}$ is an open interval containing x .

Deleted neighbourhood




Definition (Deleted neighbourhood)

A *deleted neighbourhood* of a point $x \in \mathbb{R}$ is a set formed by removing x from a neighbourhood of x .

$$(a, b) \setminus \{x\}$$

Isolated point



$$E = (a_1, b_1) \cup [a_2, b_2) \cup \{x\}$$

Definition (Isolated point)

If $x \in E \subseteq \mathbb{R}$ then x is an **isolated point** of E if there is a neighbourhood of x for which the only point in E is x itself, *i.e.*, $\exists c > 0$ such that $(x - c, x + c) \cap E = \{x\}$.

Poll

- Go to https://www.childsmath.ca/childsforms/main_login.php
- Click on [Math 3A03](#)
- Click on [Take Class Poll](#)
- Fill in poll **Lecture 13: Isolated points**
- .


Isolated point examples

Set E	Isolated points?
$(-1, 1)$	
$[0, 1]$	
\mathbb{N}	
\mathbb{R}	
\mathbb{Q}	
$(-1, 1) \cup [0, 1]$	
$(-1, 1) \setminus \{\frac{1}{2}\}$	

Isolated point examples

Set E	Isolated points?
$(-1, 1)$	\nexists
$[0, 1]$	\nexists
\mathbb{N}	Every point
\mathbb{R}	\nexists
\mathbb{Q}	\nexists
$(-1, 1) \cup [0, 1]$	\nexists
$(-1, 1) \setminus \{\frac{1}{2}\}$	\nexists

Accumulation point



$$E = \left\{ 1 - \frac{1}{n} : n \in \mathbb{N} \right\}$$

Definition (Accumulation Point or Limit Point)

If $E \subseteq \mathbb{R}$ then x is an **accumulation point** or **limit point** of E if every neighbourhood of x contains infinitely many points of E ,

$$\text{i.e.,} \quad \forall c > 0 \quad (x - c, x + c) \cap (E \setminus \{x\}) \neq \emptyset.$$

Notes:

- It is possible but not necessary that $x \in E$.
- The shorthand condition is equivalent to saying that every deleted neighbourhood of x contains at least one point of E .

Poll

- Go to https://www.childsmath.ca/childsforms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Lecture 13: Accumulation points**
- .

Accumulation point examples

Set E	Accumulation points?
$(-1, 1)$	
$[0, 1]$	
\mathbb{N}	
\mathbb{R}	
\mathbb{Q}	
$(-1, 1) \cup [0, 1]$	
$(-1, 1) \setminus \{\frac{1}{2}\}$	
$\{1 - \frac{1}{n} : n \in \mathbb{N}\}$	

Accumulation point examples

Set E	Accumulation points?
$(-1, 1)$	$[-1, 1]$
$[0, 1]$	$[0, 1]$
\mathbb{N}	\nexists
\mathbb{R}	\mathbb{R}
\mathbb{Q}	\mathbb{R}
$(-1, 1) \cup [0, 1]$	$[-1, 1]$
$(-1, 1) \setminus \{\frac{1}{2}\}$	$[-1, 1]$
$\{1 - \frac{1}{n} : n \in \mathbb{N}\}$	$\{1\}$



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 14
Topology of \mathbb{R}^n
Thursday 3 October 2019

Poll

- Go to https://www.childsmath.ca/childsa/forms/main_login.php
- Click on [Math 3A03](#)
- Click on [Take Class Poll](#)
- Fill in poll **Lecture 14: poll on polls**
- .

THINKING ABOUT GRADUATE SCHOOL?

JOIN US TO FIND OUT MORE AT THE GRAD
INFO SESSION!

WHEN: THURSDAY OCTOBER 3, 2019

TIME: 5:30PM – 7:00PM

WHERE: HH/305 AND THE MATH CAFÉ

Matheus Grasselli will give general advice on applying to grad school.

Shui Feng will talk about graduate programs particular to statistics.

Tom Hurd will talk about graduate opportunities in financial math including PhiMac.

Miroslav Lovric will give tips about applying to teachers' college.

PIZZA will be served! See you there!



Announcements

- **Assignment 3** is posted, but more problems will be added in a few days. **Due Tuesday 22 October 2019 at 2:25pm via crowdmark.**

Topological concepts covered so far

- Interval
- Neighbourhood
- Deleted neighbourhood
- Interior point
- Isolated point
- Accumulation point

Boundary point



Definition (Boundary Point)

If $E \subseteq \mathbb{R}$ then x is a **boundary point** of E if every neighbourhood of x contains at least one point of E and at least one point not in E , i.e.,

$$\forall c > 0 \quad (x - c, x + c) \cap E \neq \emptyset \\ \wedge \quad (x - c, x + c) \cap (\mathbb{R} \setminus E) \neq \emptyset.$$

Note: It is possible but not necessary that $x \in E$.

Definition (Boundary)

If $E \subseteq \mathbb{R}$ then the **boundary** of E , denoted ∂E , is the set of all boundary points of E .

Boundary point examples

Set E	Boundary points?
$(-1, 1)$	$\{-1, 1\}$
$[0, 1]$	$\{0, 1\}$
\mathbb{N}	\mathbb{N}
\mathbb{R}	\emptyset
\mathbb{Q}	\mathbb{R}
$(-1, 1) \cup [0, 1]$	$\{-1, 1\}$
$(-1, 1) \setminus \{\frac{1}{2}\}$	$\{-1, \frac{1}{2}, 1\}$
$\{1 - \frac{1}{n} : n \in \mathbb{N}\}$	$\{1 - \frac{1}{n} : n \in \mathbb{N}\} \cup \{1\}$

Closed set



Definition (Closed set)

A set $E \subseteq \mathbb{R}$ is **closed** if it contains all of its accumulation points.

Definition (Closure of a set)

If $E \subseteq \mathbb{R}$ and E' is the set of accumulation points of E then $\overline{E} = E \cup E'$ is the **closure** of E .

Note: If the set E has no accumulation points, then E is closed because there are no accumulation points to check.

Open set



Definition (Open set)

A set $E \subseteq \mathbb{R}$ is **open** if every point of E is an **interior point**.

Definition (Interior of a set)

If $E \subseteq \mathbb{R}$ then the **interior** of E , denoted $\text{int}(E)$ or E° , is the set of all **interior points** of E .

Poll

- Go to https://www.childsmath.ca/childsforms/main_login.php
- Click on [Math 3A03](#)
- Click on [Take Class Poll](#)
- Fill in poll **Lecture 14: Open or Closed**
- .

Examples

Set E	Closed?	Open?	\bar{E}	E°	∂E
$(-1, 1)$					
$[0, 1]$					
\mathbb{N}					
\mathbb{R}					
\emptyset					
\mathbb{Q}					
$(-1, 1) \cup [0, 1]$					
$(-1, 1) \setminus \{\frac{1}{2}\}$					
$\{1 - \frac{1}{n} : n \in \mathbb{N}\}$					

Examples

Set E	Closed?	Open?	\bar{E}	E°	∂E
$(-1, 1)$	NO	YES	$[-1, 1]$	E	$\{-1, 1\}$
$[0, 1]$	YES	NO	E	$(0, 1)$	$\{0, 1\}$
\mathbb{N}	YES	NO	\mathbb{N}	\emptyset	\mathbb{N}
\mathbb{R}	YES	YES	\mathbb{R}	\mathbb{R}	\emptyset
\emptyset	YES	YES	\emptyset	\emptyset	\emptyset
\mathbb{Q}	NO	NO	\mathbb{R}	\emptyset	\mathbb{R}
$(-1, 1) \cup [0, 1]$	NO	NO	$[-1, 1]$	$(-1, 1)$	$\{-1, 1\}$
$(-1, 1) \setminus \{\frac{1}{2}\}$	NO	YES	$[-1, 1]$	E	$\{-1, \frac{1}{2}, 1\}$
$\{1 - \frac{1}{n} : n \in \mathbb{N}\}$	NO	NO	$E \cup \{1\}$	\emptyset	$E \cup \{1\}$



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 15
Topology of \mathbb{R}^n III
Friday 4 October 2019

Announcements

- **Assignment 3** is posted, but more problems will be added over the weekend. **Due Tuesday 22 October 2019 at 2:25pm via [crowdmark](#).**
- **Math 3A03 Test #1**
Tuesday 29 October 2019, 5:30–7:00pm, in [JHE 264](#)
(room is booked for 90 minutes; you should not feel rushed)

Topological concepts covered so far

- Interval
- Neighbourhood
- Deleted neighbourhood
- Interior point
- Isolated point
- Accumulation point
- Boundary point
- Boundary
- Closed set
- Closure
- Open set
- Interior

Poll

- Go to https://www.childsmath.ca/childsforms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Lecture 15: The most general type of open set**
- .

Component intervals of open sets

What does the most general open set look like?

Theorem (Component intervals)

If G is an open subset of \mathbb{R} and $G \neq \emptyset$ then there is a unique (possibly finite) sequence of disjoint open intervals $\{(a_n, b_n)\}$ such that

$$G = (a_1, b_1) \cup (a_2, b_2) \cup \cdots \cup (a_n, b_n) \cup \cdots ,$$

$$\text{i.e., } G = \bigcup_{n=1}^{\infty} (a_n, b_n) .$$

The open intervals (a_n, b_n) are said to be the **component intervals** of G .

(Textbook (TBB) [Theorem 4.15](#), p. 231)

Component intervals of open sets

Main ideas of proof of [component intervals theorem](#):

- $x \in G \implies x$ is an interior point of $G \implies$
 - some neighbourhood of x is contained in G ,
i.e., $\exists c > 0$ such that $(x - c, x + c) \subseteq G$
 - \exists a largest neighbourhood of x that is contained in G : this
“**component of G** ” is $I_x = (\alpha, \beta)$, where

$$\alpha = \inf\{a : (a, x] \subset G\}, \quad \beta = \sup\{b : [x, b) \subset G\}$$

- I_x contains a rational number, i.e., $\exists r \in I_x \cap \mathbb{Q}$
- \therefore We can index all the intervals I_x by rational numbers
- \therefore There are at most countably many intervals that make up G (i.e., G is the union of a sequence of intervals)
- We can choose a disjoint subsequence of these intervals whose union is all of G (see [proof in textbook](#) for details).

Open vs. Closed Sets

Definition (Complement of a set of real numbers)

If $E \subseteq \mathbb{R}$ then the *complement* of E is the set

$$E^c = \{x \in \mathbb{R} : x \notin E\}.$$

Theorem (Open vs. Closed)

If $E \subseteq \mathbb{R}$ then E is open iff E^c is closed.

(Textbook (TBB) [Theorem 4.16](#))

Open vs. Closed Sets

Theorem (Properties of open sets of real numbers)

- 1 The sets \mathbb{R} and \emptyset are open.
- 2 Any *intersection* of a *finite* number of open sets is open.
- 3 Any *union* of an *arbitrary* collection of open sets is open.
- 4 The complement of an open set is closed.

(Textbook (TBB) [Theorem 4.17](#))

Theorem (Properties of closed sets of real numbers)

- 1 The sets \mathbb{R} and \emptyset are closed.
- 2 Any *union* of a *finite* number of closed sets is closed.
- 3 Any *intersection* of an *arbitrary* collection of closed sets is closed.
- 4 The complement of a closed set is open.

(Textbook (TBB) [Theorem 4.18](#))

Local vs. Global properties

Definition (Bounded function)


A real-valued function f is **bounded** on the set E if there exists $M > 0$ such that $|f(x)| \leq M$ for all $x \in E$.

(i.e., the function f is bounded on E iff $\{f(x) : x \in E\}$ is a bounded set.)


Note: This is a **global** property because there is a single bound M associated with the entire set E .

Example

The function $f(x) = 1/(1 + x^2)$ is bounded on \mathbb{R} . e.g., $M = 1$.




Local vs. Global properties



$f(x) = 1/x$ is not bounded on the interval $E = (0, 1)$.

Local vs. Global properties



$f(x) = 1/x$ is **locally bounded** on the interval $E = (0, 1)$,
i.e., $\forall x \in E, \exists \delta_x, M_x > 0 \mid |f(t)| \leq M_x \forall t \in (x - \delta_x, x + \delta_x)$.

Local vs. Global properties

Definition (Locally bounded at a point)

A real-valued function f is **locally bounded** at the point x if there is a neighbourhood of x in which f is bounded, *i.e.*, there exists $\delta_x > 0$ and $M_x > 0$ such that $|f(t)| \leq M_x$ for all $t \in (x - \delta_x, x + \delta_x)$.

Definition (Locally bounded on a set)

A real-valued function f is **locally bounded** on the set E if f is locally bounded at each point $x \in E$.

Note: The size of the neighbourhood (δ_x) and the local bound (M_x) depend on the point x .

Local vs. Global properties

Example (Function that is not even locally bounded)

Give an example of a function that is defined on the interval $(0, 1)$ but is not locally bounded on $(0, 1)$.

(solution on board)

Example (Function that is a mess near 0)

Give an example of a function $f(x)$ that is defined everywhere, yet in any neighbourhood of the origin there are infinitely many points at which f is not locally bounded.

(solution on board)

Extra Challenge Problem: Is there a function $f : \mathbb{R} \rightarrow \mathbb{R}$ that is not locally bounded anywhere?

Poll

- Go to https://www.childsmath.ca/childsa/forms/main_login.php
- Click on [Math 3A03](#)
- Click on [Take Class Poll](#)
- Fill in poll **Lecture 15: Local boundedness**
- .