

**13** Topology of  $\mathbb{R}$

**14** Topology of  $\mathbb{R}$  II

**15** Topology of  $\mathbb{R}$  III

**16** Topology of  $\mathbb{R}$  IV

**17** Topology of  $\mathbb{R}$  V



Mathematics  
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

# Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 13  
Topology of  $\mathbb{R}$   
Thursday 5 February 2026

# Announcements

- Solutions to [Assignment 2](#) have been posted.
- [Assignment 3](#) is posted on the course web site. Participation deadline is Tuesday 24 Feb 2026 @ 2:25pm.
- The midterm TEST is on Thursday 26 February 2026 @ 7:00pm in T13 123.
- The room is booked for 7:00–10:00 pm, but the intention is that a reasonable amount of time for the test is 90 minutes. You will be given double time.

# Topology of $\mathbb{R}$

# Intervals



*Open interval:*

$$(a, b) = \{x : a < x < b\}$$


*Closed interval:*

$$[c, d] = \{x : c \leq x \leq d\}$$

*Half-open interval:*

$$(e, f] = \{x : e < x \leq f\}$$

# Interior point



## Definition (Interior point)


If  $E \subseteq \mathbb{R}$  then  $x$  is an *interior point* of  $E$  if  $x$  lies in an open interval that is contained in  $E$ , i.e.,

$$\exists c > 0 \quad \text{) } (x - c, x + c) \subset E.$$

## Interior point examples

Set $E$	Interior points?
$(-1, 1)$	
$[0, 1]$	
$\mathbb{N}$	
$\mathbb{R}$	
$\mathbb{Q}$	
$(-1, 1) \cup [0, 1]$	
$(-1, 1) \setminus \{\frac{1}{2}\}$	


# Neighbourhood



## Definition (Neighbourhood)

A **neighbourhood** of a point  $x \in \mathbb{R}$  is an open interval containing  $x$ .

# Deleted neighbourhood



$$(a, b) \setminus \{x\}$$

## Definition (Deleted neighbourhood)

A *deleted neighbourhood* of a point  $x \in \mathbb{R}$  is a set formed by removing  $x$  from a neighbourhood of  $x$ .

# Isolated point



$$E = (a_1, b_1) \cup [a_2, b_2) \cup \{x\}$$

## Definition (Isolated point)

If  $x \in E \subseteq \mathbb{R}$  then  $x$  is an **isolated point** of  $E$  if there is a neighbourhood of  $x$  for which the only point in  $E$  is  $x$  itself, *i.e.*,

$$\exists c > 0 \quad \} \quad (x - c, x + c) \cap E = \{x\}.$$

# Poll

- Go to  
[https://www.childsmath.ca/childsa/forms/main\\_login.php](https://www.childsmath.ca/childsa/forms/main_login.php)
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Topology: Isolated points**
- .

## Isolated point examples

Set $E$	Isolated points?
$(-1, 1)$	
$[0, 1]$	
$\mathbb{N}$	
$\mathbb{R}$	
$\mathbb{Q}$	
$(-1, 1) \cup [0, 1]$	
$(-1, 1) \setminus \{\frac{1}{2}\}$	

# Accumulation point



$$E = \left\{ 1 - \frac{1}{n} : n \in \mathbb{N} \right\}$$

## Definition (Accumulation Point or Limit Point or Cluster Point)

If  $E \subseteq \mathbb{R}$  then  $x$  is an **accumulation point** of  $E$  if every neighbourhood of  $x$  contains infinitely many points of  $E$ ,

$$\text{i.e., } \forall c > 0 \quad (x - c, x + c) \cap (E \setminus \{x\}) \neq \emptyset.$$

### Note:

- It is possible but not necessary that  $x \in E$ .
- The shorthand condition is equivalent to saying that every deleted neighbourhood of  $x$  contains at least one point of  $E$ .

# Poll

- Go to  
[https://www.childsmath.ca/childsa/forms/main\\_login.php](https://www.childsmath.ca/childsa/forms/main_login.php)
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Topology: Accumulation points**
- .

## Accumulation point examples

Set $E$	Accumulation points?
$(-1, 1)$	
$[0, 1]$	
$\mathbb{N}$	
$\mathbb{R}$	
$\mathbb{Q}$	
$(-1, 1) \cup [0, 1]$	
$(-1, 1) \setminus \{\frac{1}{2}\}$	
$\{1 - \frac{1}{n} : n \in \mathbb{N}\}$	

# Boundary point



## Definition (Boundary Point)

If  $E \subseteq \mathbb{R}$  then  $x$  is a **boundary point** of  $E$  if every neighbourhood of  $x$  contains at least one point of  $E$  and at least one point not in  $E$ , i.e.,

$$\forall c > 0 \quad \begin{aligned} (x - c, x + c) \cap E &\neq \emptyset \\ \wedge \quad (x - c, x + c) \cap (\mathbb{R} \setminus E) &\neq \emptyset. \end{aligned}$$

Note: It is possible but not necessary that  $x \in E$ .

## Definition (Boundary)

If  $E \subseteq \mathbb{R}$  then the **boundary** of  $E$ , denoted  $\partial E$ , is the set of all boundary points of  $E$ .

## Boundary point examples

Set $E$	Boundary points?
$(-1, 1)$	
$[0, 1]$	
$\mathbb{N}$	
$\mathbb{R}$	
$\mathbb{Q}$	
$(-1, 1) \cup [0, 1]$	
$(-1, 1) \setminus \{\frac{1}{2}\}$	
$\{1 - \frac{1}{n} : n \in \mathbb{N}\}$	

# Closed set



## Definition (Closed set)

A set  $E \subseteq \mathbb{R}$  is **closed** if it contains all of its accumulation points.

## Definition (Closure of a set)

If  $E \subseteq \mathbb{R}$  and  $E'$  is the set of accumulation points of  $E$  then the **closure** of  $E$  is

$$\bar{E} = E \cup E'.$$

**Note:** If the set  $E$  has no accumulation points, then  $E$  is closed because there are no accumulation points to check.



# Poll

- Go to  
[https://www.childsmath.ca/childsa/forms/main\\_login.php](https://www.childsmath.ca/childsa/forms/main_login.php)
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Topology: Open or Closed**
- .

## Examples

Set $E$	Closed?	Open?	$\bar{E}$	$E^\circ$	$\partial E$
$(-1, 1)$					
$[0, 1]$					
$\mathbb{N}$					
$\mathbb{R}$					
$\emptyset$					
$\mathbb{Q}$					
$(-1, 1) \cup [0, 1]$					
$(-1, 1) \setminus \{\frac{1}{2}\}$					
$\{1 - \frac{1}{n} : n \in \mathbb{N}\}$					



Mathematics  
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

# Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 14  
Topology of  $\mathbb{R}^n$   
Friday 6 February 2026

# Announcements

- Today's Mathematics Colloquium (immediately after this class in HH-305) should be accessible to 3A students. Title "[Is mathematics obsolete?](#)"
- Solutions to [Assignment 2](#) have been posted.
- [Assignment 3](#) is posted on the course web site. Participation deadline is Tuesday 24 Feb 2025 @ 2:25pm.
- The midterm TEST is on Thursday 26 February 2026 @ 7:00pm in T13 123.
- The room is booked for 7:00–10:00 pm, but the intention is that a reasonable amount of time for the test is 90 minutes. You will be given double time.
- Last year's (winter 2025) midterm test and solutions are posted on the [course web site](#).

# Topological concepts covered so far

- Interval
- Neighbourhood
- Deleted neighbourhood
- Interior point
- Isolated point
- Accumulation point
- Boundary point
- Boundary
- Closed set
- Closure
- Open set
- Interior

# Equivalent definitions

## Example (Closure)

For  $E \subseteq \mathbb{R}$ , prove  $E \cup E' = E \cup \partial E$ , so  $\bar{E}$  can be defined either way.

*Proof:* We must show  $E \cup E' \subseteq E \cup \partial E$  and  $E \cup E' \supseteq E \cup \partial E$ .

- $\subseteq$  Suppose  $x \in E \cup E'$ . If  $x \in E$  then  $x \in E \cup A$  for any set  $A$ . In particular,  $x \in E \cup \partial E$ . Alternatively, suppose  $x \notin E$ , i.e.,  $x \in E^c$ . Then, since  $x \in E \cup E'$ , it must be that  $x \in E'$ , which means that any neighbourhood of  $x$  contains a point of  $E$ . But  $x \in E^c$ , so any such neighbourhood also contains a point of  $E^c$  (namely  $x$ ). Therefore,  $x \in \partial E \subseteq E \cup \partial E$ .
- $\supseteq$  Suppose  $x \in E \cup \partial E$ . If  $x \in E$  then  $x \in E \cup A$  for any set  $A$ . In particular,  $x \in E \cup E'$ . Alternatively, suppose  $x \notin E$ , i.e.,  $x \in E^c$ . Then, since  $x \in E \cup \partial E$ , it must be that  $x \in \partial E$ , which means that any neighbourhood of  $x$  contains a point of  $E$ . But that point is not  $x$ , since  $x \notin E$ . Thus, any *deleted* neighbourhood of  $x$  contains a point of  $E$ , i.e.,  $x \in E' \subseteq E \cup E'$ . □

*Question:* In the proof above, did we use any properties of  $\mathbb{R}$  ?

# Poll

- Go to  
[https://www.childsmath.ca/childsa/forms/main\\_login.php](https://www.childsmath.ca/childsa/forms/main_login.php)
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Topology: The most general type of open set**
- .

# Component intervals of open sets

What does the most general **open set** look like?

## Theorem (Component intervals)

If  $G$  is an open subset of  $\mathbb{R}$  and  $G \neq \emptyset$  then there is a unique (possibly finite) sequence of disjoint open intervals  $\{(a_n, b_n)\}$  such that

$$G = (a_1, b_1) \cup (a_2, b_2) \cup \cdots \cup (a_n, b_n) \cup \cdots ,$$

$$\text{i.e., } G = \bigcup_{n=1}^{\infty} (a_n, b_n) .$$

The open intervals  $(a_n, b_n)$  are said to be the **component intervals** of  $G$ .

(TBB [Theorem 4.15](#), p. 231)

# Component intervals of open sets

Main ideas of proof of **component intervals theorem**:

- $x \in G \implies x \in G^\circ \implies x \in (a, b) \subset G$  for some  $a, b \in \mathbb{R}$
- $\exists$  a largest neighbourhood of  $x$  in  $G$ : this **component of  $G$**  is  $I_x = (\alpha, \beta)$ , where  $\alpha = \inf\{a : (a, x] \subset G\}$  and  $\beta = \sup\{b : [x, b) \subset G\}$
- Components with different endpoints cannot overlap (they would contradict the inf and sup) so **distinct components are disjoint**
- Every component  $I_x$  **contains a rational number**, i.e.,  $\exists r \in I_x \cap \mathbb{Q}$
- But  $\mathbb{Q}$  is **countable**, and we can write an **explicit bijection**  $q : \mathbb{N} \rightarrow \mathbb{Q}$
- $\therefore$  We can **label** each component interval  $I$  of  $G$  with a **unique**  $n \in \mathbb{N}$ . The label of component  $I$  is  $n_I = \min\{n \in \mathbb{N} : q(n) \in I\}$  (well-ordering)
- $\therefore$  We can list the components of  $G$  in a **sequence**  $\{I_1, I_2, I_3, \dots\}$
- $\therefore$  There are at most countably many intervals that make up  $G$  i.e.,  **$G$  is the union of a sequence of disjoint intervals**
- See **textbook** for a standard proof (TBB **Theorem 4.15**, p. 231).

Note: (For those studying logic and set theory.) This argument relies ultimately on the **well-ordering principle for  $\mathbb{N}$**  (or, equivalently, the principle of mathematical induction for  $\mathbb{N}$ ) but it does not require the **Axiom of Choice**.

# Open vs. Closed Sets

## Definition (Complement of a set of real numbers)

If  $E \subseteq \mathbb{R}$  then the **complement** of  $E$  is the set

$$E^c = \{x \in \mathbb{R} : x \notin E\}.$$

## Theorem (Open vs. Closed)

*If  $E \subseteq \mathbb{R}$  then  $E$  is open iff  $E^c$  is closed.*

(TBB [Theorem 4.16](#))

# Open vs. Closed Sets

## Theorem (Properties of open sets of real numbers)

- 1 The sets  $\mathbb{R}$  and  $\emptyset$  are open.
- 2 Any *intersection* of a *finite* number of open sets is open.
- 3 Any *union* of an *arbitrary* collection of open sets is open.
- 4 The complement of an open set is closed.

(TBB Theorem 4.17)

## Theorem (Properties of closed sets of real numbers)

- 1 The sets  $\mathbb{R}$  and  $\emptyset$  are closed.
- 2 Any *union* of a *finite* number of closed sets is closed.
- 3 Any *intersection* of an *arbitrary* collection of closed sets is closed.
- 4 The complement of a closed set is open.

(TBB Theorem 4.18)



Mathematics  
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

# Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 15  
Topology of  $\mathbb{R}^n$   
Tuesday 10 February 2026

# Announcements

- Solutions to [Assignment 2](#) have been re-posted (some typos have been fixed).
- [Assignment 3](#) is posted on the course web site. Participation deadline is Tuesday 24 Feb 2025 @ 2:25pm.
- The midterm TEST is on Thursday 26 February 2026 @ 7:00pm in T13 123.
- The room is booked for 7:00–10:00 pm, but the intention is that a reasonable amount of time for the test is 90 minutes. You will be given double time.
- Last year's (winter 2025) midterm test and solutions are posted on the [course web site](#).
- I have revised my [sketch](#) of the proof of the [component intervals theorem](#).
- For those who are interested in foundations of mathematics: the simplest result in real analysis that depends on the Axiom of Choice is the existence of a non-measurable set (take Math 4A...)

# Local vs. Global properties

## Definition (Bounded function)


A real-valued function  $f$  is **bounded** on the set  $E$  if there exists  $M > 0$  such that  $|f(x)| \leq M$  for all  $x \in E$ .

(i.e., the function  $f$  is bounded on  $E$  iff  $\{f(x) : x \in E\}$  is a bounded set.)


Note: This is a **global** property because there is a single bound  $M$  associated with the entire set  $E$ .

## Example

The function  $f(x) = 1/(1 + x^2)$  is bounded on  $\mathbb{R}$ . e.g.,  $M = 1$ .




# Local vs. Global properties



$f(x) = 1/x$  is not bounded on the interval  $E = (0, 1)$ .

## Local vs. Global properties



$f(x) = 1/x$  is **locally bounded** on the interval  $E = (0, 1)$ ,  
i.e.,  $\forall x \in E, \exists \delta_x, M_x > 0 \mid |f(t)| \leq M_x \forall t \in (x - \delta_x, x + \delta_x)$ .

# Local vs. Global properties

## Definition (Locally bounded at a point)

A real-valued function  $f$  is **locally bounded** at the point  $x$  if there is a neighbourhood of  $x$  in which  $f$  is bounded, *i.e.*, there exists  $\delta_x > 0$  and  $M_x > 0$  such that  $|f(t)| \leq M_x$  for all  $t \in (x - \delta_x, x + \delta_x)$ .

## Definition (Locally bounded on a set)

A real-valued function  $f$  is **locally bounded** on the set  $E$  if  $f$  is locally bounded at each point  $x \in E$ .

Note: The size of the neighbourhood ( $\delta_x$ ) and the local bound ( $M_x$ ) depend on the point  $x$ .

# Local vs. Global properties

## Example (Function that is not even locally bounded)

Give an example of a function that is defined on the interval  $(0, 1)$  but is not **locally bounded** on  $(0, 1)$ .

Let's construct a function  $f(x)$  that is defined on  $(0, 1)$  but is not locally bounded at one point, say  $x = \frac{1}{2}$ .

$f(x)$  must blow up  $x = \frac{1}{2}$ . Let's make  $f$  look like  $1/x$ , but shifted so the blowup is at  $x = \frac{1}{2}$ .

$$f(x) = \begin{cases} \frac{1}{x - \frac{1}{2}} & x \neq \frac{1}{2}, \\ 0 & x = \frac{1}{2}. \end{cases}$$

# Local vs. Global properties

## Example (Function that is a mess near 0)

Give an example of a function  $f(x)$  that is defined everywhere, yet in any neighbourhood of the origin there are infinitely many points at which  $f$  is not locally bounded.

**Please do poll: Topology: Local boundedness**

$$\text{Consider } S(x) = \begin{cases} \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

$S(x)$  is bounded on  $\mathbb{R}$ , hence locally bounded at every point.

$$\text{Consider } T(x) = \begin{cases} \tan \frac{1}{x}, & x \neq 0 \text{ and } \cos \frac{1}{x} \neq 0 \\ 0, & x = 0 \text{ or } \cos \frac{1}{x} = 0 \end{cases}$$

$T(x)$  is not locally bounded at points where  $\cos \frac{1}{x} = 0$ , i.e., for  $\frac{1}{x} = \frac{\pi}{2} + n\pi$ ,  $n \in \mathbb{Z}$ . There are infinitely many such points in any neighbourhood of  $x = 0$ .

# Poll

- Go to  
[https://www.childsmath.ca/childsa/forms/main\\_login.php](https://www.childsmath.ca/childsa/forms/main_login.php)
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Topology: Locally bounded nowhere?**
- .

# Local vs. Global properties

## Extra Challenge Problem:

Is there a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  that is not locally bounded anywhere?

# Local vs. Global properties

- What condition(s) rule out such pathological behaviour?
- When does a property holding locally (near any given point in a set) imply that it holds globally (for the set as a whole)?
- For example: What condition(s) must a set  $E \subseteq \mathbb{R}$  satisfy in order that a function  $f$  that is **locally bounded** on  $E$  is necessarily **bounded** on  $E$ ?
- We will see that the condition we are seeking is that the set  $E$  must be “**compact**” ...

# Compactness

Recall the Bolzano-Weierstrass theorem, for sequences of real numbers:

Theorem (Bolzano-Weierstrass theorem for sequences)

*Every bounded sequence in  $\mathbb{R}$  contains a convergent subsequence.*

For *any set of real numbers*, we define:

Definition (Bolzano-Weierstrass property)

A set  $E \subseteq \mathbb{R}$  is said to have the ***Bolzano-Weierstrass property*** iff any sequence of points chosen from  $E$  has a subsequence that converges to a point in  $E$ .

# Compactness

## Theorem (Bolzano-Weierstrass theorem for sets)

A set  $E \subseteq \mathbb{R}$  has the *Bolzano-Weierstrass property* iff  $E$  is closed and bounded.

(TBB [Theorem 4.21](#), p. 241)

## Proof of $\Leftarrow$ .

Suppose  $E$  is closed and bounded. Let  $\{x_n\}$  be a sequence in  $E$ . Since  $E$  is bounded, the usual Bolzano-Weierstrass theorem implies that there is a subsequence  $\{x_{n_k}\}$  that converges. If  $\{x_{n_k}\}$  is eventually constant, then its limit is a point in  $E$ , so we're done. Otherwise,  $\{x_{n_k}\}$  must converge to an accumulation point of  $E$ . But  $E$  is closed, so it contains all its accumulation points, including the limit of  $\{x_{n_k}\}$ . Thus, again, we have a subsequence of  $\{x_n\}$  that converges to a point in  $E$ .  $\square$

# Compactness

## Theorem (Bolzano-Weierstrass theorem for sets)

A set  $E \subseteq \mathbb{R}$  has the *Bolzano-Weierstrass property* iff  $E$  is closed and bounded.

### Proof of $\implies$ (Part 1)

Let's prove the contrapositive, *i.e.*, If  $E$  is either not bounded or not closed then  $E$  does not have the Bolzano-Weierstrass property. **Suppose  $E$  is unbounded.** In particular, suppose  $E$  is not bounded above (the argument is similar if  $E$  is not bounded below). We will construct a sequence in  $E$  that has no convergent subsequence. Pick a point  $x_1 \in E$  such that  $x_1 > 1$  (which is possible because  $E$  is not bounded above). Also, since  $E$  is not bounded above, we can find  $x_2 \in E$  such that  $x_2 > x_1 + 1 > 2$ . More generally, given  $x_k \in E$  we can find  $x_{k+1} \in E$  such that  $x_{k+1} > x_k + 1 > k + 1$ . The sequence  $\{x_n\}$  constructed in this way is increasing and diverges to  $\infty$  (since  $x_n > n$  for all  $n \in \mathbb{N}$ ). Moreover, this is true of any subsequence of  $\{x_n\}$ .  $\therefore$   **$E$  does not have the Bolzano-Weierstrass property.**

# Compactness

## Theorem (Bolzano-Weierstrass theorem for sets)

A set  $E \subseteq \mathbb{R}$  has the *Bolzano-Weierstrass property* iff  $E$  is closed and bounded.

### Proof of $\implies$ (Part 2)

Now *suppose  $E$  is not closed*. Then there must be a sequence  $\{x_n\}$  in  $E$  such that  $\{x_n\}$  converges to a point not in  $E$ . If  $E$  has the Bolzano-Weierstrass property, then  $\{x_n\}$  has a subsequence that converges to a point in  $E$ . But every subsequence of a convergent sequence must converge to the same point as the full sequence, and the full sequence converges to a point not in  $E$ ! Thus, if  $E$  is not closed *then it does not have the Bolzano-Weierstrass property*. □



Mathematics  
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

# Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 16  
Topology of  $\mathbb{R}^n$  IV  
Thursday 12 February 2026

# Announcements

- The participation deadline for [Assignment 3](#) is Tuesday 24 Feb 2025 @ 2:25pm.
- The midterm TEST is on Thursday 26 February 2026 @ 7:00pm in T13 123.
- The room is booked for 7:00–10:00 pm, but the intention is that a reasonable amount of time for the test is 90 minutes. You will be given double time.
- Last year's (winter 2025) midterm test and solutions are posted on the [course web site](#).

# Compactness

## Theorem (Bolzano-Weierstrass theorem for sets)

A set  $E \subseteq \mathbb{R}$  has the *Bolzano-Weierstrass property* iff  $E$  is closed and bounded.

### Notes:

- Why do we need both *closed* and *bounded*? Why don't we need *closed* in the *Bolzano-Weierstrass theorem* for sequences?
  - Because the statement of the *Bolzano-Weierstrass theorem* for sequences doesn't require the limit of the convergent subsequence to be in the set!
- The *Bolzano-Weierstrass theorem for sets* implies that "If  $E \subseteq \mathbb{R}$  is bounded then its closure  $\overline{E}$  has the Bolzano-Weierstrass property".
  - The *Bolzano-Weierstrass theorem for sequences* is a special case of this statement because any convergent sequence together with its limit is a closed set.

# Compactness

## Definition (Open Cover)

Let  $E \subseteq \mathbb{R}$  and let  $\mathcal{U}$  be a family of open intervals. If for every  $x \in E$  there exists at least one interval  $U \in \mathcal{U}$  such that  $x \in U$ , i.e.,

$$E \subseteq \bigcup \{U : U \in \mathcal{U}\},$$

then  $\mathcal{U}$  is called an *open cover* of  $E$ .

## Example (Open covers of $\mathbb{N}$ )

Give examples of open covers of  $\mathbb{N}$ .

- $\mathcal{U} = \left\{ \left( n - \frac{1}{2}, n + \frac{1}{2} \right) : n = 1, 2, \dots \right\}$
- $\mathcal{U} = \{(0, \infty)\}$
- $\mathcal{U} = \{(0, \infty), \mathbb{R}, (\pi, 27)\}$

# Compactness

Example (Open covers of  $\{\frac{1}{n} : n \in \mathbb{N}\}$ )

- $\mathcal{U} = \{(0, 1), (0, 2), \mathbb{R}, (\pi, 27)\}$
- $\mathcal{U} = \{(0, 2)\}$
- $\mathcal{U} = \left\{ \left( \frac{1}{n}, \frac{1}{n} + \frac{3}{4} \right) : n = 1, 2, \dots \right\}$

Example (Open covers of  $[0, 1]$ )

- $\mathcal{U} = \{(-2, 2)\}$
- $\mathcal{U} = \left\{ \left( -\frac{1}{2}, \frac{1}{2} \right), (0, 2) \right\}$
- $\mathcal{U} = \left\{ \left( \frac{1}{n}, 2 \right) : n = 1, 2, \dots \right\} \cup \left\{ \left( -\frac{1}{2}, \frac{1}{2} \right) \right\}$

# Poll

- Go to  
[https://www.childsmath.ca/childsa/forms/main\\_login.php](https://www.childsmath.ca/childsa/forms/main_login.php)
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Topology: Open covers of inverse squares**
- .

# Compactness

## Definition (Heine-Borel Property)

A set  $E \subseteq \mathbb{R}$  is said to have the *Heine-Borel property* if every open cover of  $E$  can be reduced to a finite subcover. That is, if  $\mathcal{U}$  is an open cover of  $E$ , then there exists a finite subfamily  $\{U_1, U_2, \dots, U_n\} \subseteq \mathcal{U}$ , such that  $E \subseteq U_1 \cup U_2 \cup \dots \cup U_n$ .

When does any open cover of a set  $E$  have a finite subcover?

## Theorem (Heine-Borel Theorem)

A set  $E \subseteq \mathbb{R}$  has the *Heine-Borel property* iff  $E$  is both *closed* and *bounded*.

(TBB pp. 249–250)

# Compactness

## Definition (Compact Set)

A set  $E \subseteq \mathbb{R}$  is said to be **compact** if it has any of the following equivalent properties:

- 1  $E$  is **closed** and bounded.
- 2  $E$  has the **Bolzano-Weierstrass property**.
- 3  $E$  has the **Heine-Borel property**.

Note: In spaces other than  $\mathbb{R}$ , these three properties are not necessarily equivalent. Usually the **Heine-Borel property** is taken as the definition of compactness.

# Compactness

## Example

Prove that the interval  $(0, 1]$  is not compact by showing that it is not closed or not bounded.

## Example

Prove that the interval  $(0, 1]$  is not compact by showing that it does not have the Bolzano-Weierstrass property.

## Example

Prove that the interval  $(0, 1]$  is not compact by showing that it does not have the Heine-Borel property.

# Poll

- Go to  
[https://www.childsmath.ca/childsa/forms/main\\_login.php](https://www.childsmath.ca/childsa/forms/main_login.php)
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Topology: Compactness**
- .

# Topological concepts that we have covered

- Interval
- Neighbourhood
- Deleted neighbourhood
- Interior point
- Isolated point
- Accumulation point
- Boundary point
- Boundary
- Closed set
- Closure
- Open set
- Interior
- Complement
- Compact

# Compactness

Classic non-trivial compactness argument:

Theorem (Compact  $\implies$  bounded if locally bounded)

Let  $E$  be a *compact* subset of  $\mathbb{R}$ . If  $f : E \rightarrow \mathbb{R}$  is *locally bounded* on  $E$  then  $f$  is *bounded* on  $E$ .

Proof via Bolzano-Weierstrass.

Suppose  $f$  is *locally bounded* on  $E$ , and that  $E$  satisfies the *Bolzano-Weierstrass property*. Suppose further, in order to derive a contradiction, that  $f$  is not bounded on  $E$ . Then there is some sequence  $\{x_n\}$  in  $E$  such that  $|f(x_n)| > n$  (otherwise  $|f(x_n)| \leq N$  for some  $N \in \mathbb{N}$ , so  $N$  would be a bound). By the Bolzano-Weierstrass property,  $\{x_n\}$  has a subsequence  $\{x_{n_k}\}$  such that  $x_{n_k} \rightarrow L \in E$ . Since  $f$  is locally bounded, there exist  $\delta > 0$  and  $M > 0$  such that  $|f(x)| \leq M$  for all  $x \in (L - \delta, L + \delta)$ . But  $x_{n_k} \rightarrow L$ , so for all sufficiently large  $k$ ,  $x_{n_k} \in (L - \delta, L + \delta)$ . Yet for sufficiently large  $k$ ,  $|f(x_{n_k})| > n_k \geq k > M$ .  $\implies \Leftarrow$ . Hence  $f$  must, in fact, be bounded on  $E$ .  $\square$

# Compactness

Theorem (Compact  $\implies$  bounded if locally bounded)

Let  $E$  be a *compact* subset of  $\mathbb{R}$ . If  $f : E \rightarrow \mathbb{R}$  is *locally bounded* on  $E$  then  $f$  is *bounded* on  $E$ .

Proof via Heine-Borel.

Since  $f$  is locally bounded, each  $x \in E$  lies in some open interval  $U_x$  such that  $|f(t)| \leq M_x$  for all  $t \in U_x$ . The collection  $\mathcal{U} = \{U_x : x \in E\}$  is an open cover of  $E$ . But  $E$  satisfies the *Heine-Borel property*, so  $\mathcal{U}$  contains a finite subcover, say  $\{U_{x_1}, \dots, U_{x_n}\}$ . We can therefore find a bound for  $f$  on all of  $E$ . Let  $M = \max\{M_{x_1}, \dots, M_{x_n}\}$ . Then  $|f(x)| \leq M$  for all  $x \in E$ .  $\square$

Proof via Heine-Borel is much easier!



Mathematics  
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

# Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 17  
Topology of  $\mathbb{R}^n$   
Friday 13 February 2026

# Announcements

- The participation deadline for [Assignment 3](#) is Tuesday 24 Feb 2025 @ 2:25pm.
- The midterm TEST is on Thursday 26 February 2026 @ 7:00pm in T13 123.
- The room is booked for 7:00–10:00 pm, but the intention is that a reasonable amount of time for the test is 90 minutes. You will be given double time.
- Last year's (winter 2025) midterm test and solutions are posted on the [course web site](#).

# Compactness

Theorem (Compact  $\implies$  bounded if locally bounded)

Let  $E$  be a *compact* subset of  $\mathbb{R}$ . If  $f : E \rightarrow \mathbb{R}$  is *locally bounded* on  $E$  then  $f$  is *bounded* on  $E$ .

Example (Converse of above theorem)

Let  $E \subseteq \mathbb{R}$ . If every function  $f : E \rightarrow \mathbb{R}$  that is *locally bounded* on  $E$  is *bounded* on  $E$ , then  $E$  is *compact*.

Note: The contrapositive of the converse is: If  $E \subseteq \mathbb{R}$  is not compact then  $\exists f : E \rightarrow \mathbb{R}$   $\nexists$   $f$  is *locally bounded* on  $E$  but not bounded on  $E$ .

# Compactness

## Example (“bounded if locally bounded” $\implies$ compact)

Consider the contrapositive: if  $E \subseteq \mathbb{R}$  is not compact then there exists a function  $f : E \rightarrow \mathbb{R}$  that is locally bounded but not bounded on  $E$ .

Suppose  $E$  is not compact. Then either  $E$  is not bounded or not closed.

Suppose first that  $E$  is not bounded, and let  $f(x) = x$ . Then  $f$  is locally bounded at any point  $x \in E$ , since  $f$  is bounded on any neighbourhood of  $x$  that has finite width; but  $f(E) = E$  is an unbounded set, so  $f$  is an unbounded function on  $E$ .

Now suppose  $E$  is not closed. Then there is an accumulation point of  $E$  that is not in  $E$ , i.e., there exists a sequence  $\{x_n\} \subseteq E$  such that  $x_n \rightarrow L \notin E$  as  $n \rightarrow \infty$ . Since  $L \notin E$ , for any  $x \in E$  we have  $x - L \neq 0$ , so we can define  $f(x) = \frac{1}{x-L}$  for all  $x \in E$ . But since  $L$  is a limit point of  $E$ , there are points in  $E$  that are arbitrarily close to  $L$ , hence points at which  $|f(x)|$  is arbitrarily large. Thus,  $f$  is unbounded on  $E$ .  $\square$

# Complements and Closures problem

## Example

How many distinct sets can be obtained from  $E = [0, 1]$  by applying the complement and closure operations?

Consider this sequence of sets:

$$E_1 = [0, 1],$$

$$E_2 = E_1^c = (-\infty, 0) \cup (1, \infty),$$

$$E_3 = \overline{E_2} = (-\infty, 0] \cup [1, \infty),$$

$$E_4 = E_3^c = (0, 1),$$

$$E_5 = \overline{E_4} = E_1.$$

Is the answer 4 for any set  $E \subseteq \mathbb{R}$ ?

# Poll

- Go to  
[https://www.childsmath.ca/childsa/forms/main\\_login.php](https://www.childsmath.ca/childsa/forms/main_login.php)
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Topology: Complements and Closures**
- .

# Complements and Closures problem

## Extra Challenge Problem

If  $E \subseteq \mathbb{R}$ , how many distinct sets can be obtained by taking complements or closures of  $E$  and its successors? Put another way, if  $\{E_n\}$  is a sequence of sets produced by taking the complement or closure of the previous set, how many distinct sets can such a sequence contain? If the answer is finite, find a set  $E$  that generates the maximum number in this way.

# Midterm Test

*What you need to know:*

- Everything discussed in class, including all definitions/concepts and theorems/lemmas/corollaries.
- Everything in assignments and solutions to assignments.  
*Make sure you fully understand all the solutions to all the problems in all the assignments.*
- Most—but not all—of the material that you are responsible for is covered in the chapters of the textbooks indicated on the course web page. You are not responsible for material in the textbooks that was not mentioned in lectures, tutorials or assignments.
- It is essential that you understand how to use the definitions and theorems to construct proofs.