

19 Sequences and Series of Functions

20 Sequences and Series of Functions II

21 Sequences and Series of Functions III

22 Sequences and Series of Functions IV

Poll

- Go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Post-Test**
- .



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 19
Sequences and Series of Functions
Friday 27 February 2026

Announcements

- New, exciting topic today...
- Assignment 4 is posted on the course web site.

Sequences and Series of Functions

Limits of Functions

We know that it can be useful to represent functions as limits of other functions.

Example

The power series expansion

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots$$

expresses the exponential e^x as a certain limit of the functions

$$1, \quad 1 + \frac{x}{1!}, \quad 1 + \frac{x}{1!} + \frac{x^2}{2!}, \quad 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!}, \quad \cdots$$

Our goal is to give meaning to the phrase “*limit of functions*”, and discuss how functions behave under limits.

Pointwise Convergence

- There are multiple inequivalent ways to define the limit of a sequence of functions.
- Consequently, there are multiple different notions of what it means for a sequence of functions to converge.
- Some convergence notions are better behaved than others.

We will begin with the simplest notion of convergence.

Definition (Pointwise Convergence)

Suppose $\{f_n\}$ is a sequence of functions defined on a domain $D \subseteq \mathbb{R}$, and let f be another function defined on D . Then $\{f_n\}$ **converges pointwise on D to f** if, for every $x \in D$, the sequence $\{f_n(x)\}$ of real numbers converges to $f(x)$.

What useful properties of functions does *pointwise convergence* preserve?


Poll

- Go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Function sequences: Pointwise convergence**
- .

Pointwise Convergence

Example

$$f_n(x) = \begin{cases} x^n & 0 \leq x \leq 1, \\ 1 & x \geq 1. \end{cases}$$



$$\lim_{n \rightarrow \infty} f_n(x) = \begin{cases} 0 & 0 \leq x < 1 \\ 1 & x \geq 1 \end{cases}$$


- The limit of this sequence (of continuous functions) is not continuous.
- If we smooth the corner of $f_n(x)$ at $x = 1$, we get a sequence of differentiable functions that converge to a function that is not even continuous.

Pointwise Convergence

Example

Define $f_n(x)$ on $[0, 1]$ as follows:

$$f_n(x) = \begin{cases} 2n^2x, & 0 \leq x \leq \frac{1}{2n} \\ 2n - 2n^2x, & \frac{1}{2n} \leq x \leq \frac{1}{n} \\ 0, & x \geq \frac{1}{n}. \end{cases}$$



$$\lim_{n \rightarrow \infty} f_n(x) = 0 \quad \forall x$$

$$\int_0^1 f_n = \frac{1}{2} \quad \forall n \in \mathbb{N}$$

$$\int_0^1 \lim_{n \rightarrow \infty} f_n = 0$$

Pointwise Convergence

In the [previous example](#), each f_n is integrable and the limit function (the zero function) is also integrable. The example shows that, nevertheless, the sequence of integrals $\{\int f_n\}$ need not converge to the integral of the limit function $\int f$.

Is pointwise convergence sufficient for integrability to be passed on to the limit function?

If so, how do we prove it?

If not, what is a counter-example?


Pointwise Convergence

Example

Let's try to construct a sequence of functions that converges to a non-integrable function. One approach is for each f_n to be bounded, but to converge to an unbounded function.

$$f_n(x) = \begin{cases} 0 & x < \frac{1}{n}, \\ \frac{1}{x} & \frac{1}{n} < x \leq 1, \\ 0 & 1 < x. \end{cases}$$

$$\lim_{n \rightarrow \infty} f_n(x) = \begin{cases} 0 & x \leq 0 \\ \frac{1}{x} & 0 < x \leq 1 \\ 0 & 1 < x \end{cases}$$



- The limit of this sequence (of integrable functions) is not bounded on $[0, 1]$, hence not integrable on $[0, 1]$.
- What if $|f_n(x)| \leq M \forall x \forall n$, i.e., if $\{f_n\}$ is **uniformly bounded**?

Pointwise Convergence

Example

Let's try to construct a *uniformly bounded* sequence of functions that converges to a non-integrable, bounded function. Recall the non-integrable function

$$f(x) = \begin{cases} 1 & x \in \mathbb{Q}, \\ 0 & x \notin \mathbb{Q}. \end{cases}$$

Let's construct a sequence of integrable functions that converges to f . Since $\mathbb{Q}$ is countable, we can list all of its elements in a sequence $\{q_k : k = 1, 2, \dots\}$. Now define f_n on $[0, 1]$ via

$$f_n(x) = \begin{cases} 1 & x \in \{q_1, \dots, q_n\}, \\ 0 & \text{otherwise.} \end{cases}$$

Then $f_n(x) \leq 1 \ \forall x$ and on any closed interval (e.g., $[0, 1]$) each f_n is integrable, since it is piecewise continuous, but $f_n \rightarrow f$, which is not integrable on any interval. □

Uniform Convergence

A much better behaved notion of convergence is the following.

Definition ($f_n \rightarrow f$ uniformly)

Suppose $\{f_n\}$ is a sequence of functions defined on a domain $D \subseteq \mathbb{R}$, and let f be another function defined on D . Then $\{f_n\}$ **converges uniformly on D to f** if, for every $\varepsilon > 0$, there is some $N \in \mathbb{N}$ so that, for all $x \in D$,

$$n \geq N \implies |f_n(x) - f(x)| < \varepsilon.$$

Note that $\{f_n\}$ **converges uniformly** to f if and only if $\forall \varepsilon > 0$, $\exists N \in \mathbb{N}$ such that

$$n \geq N \implies \sup_{x \in D} |f_n(x) - f(x)| < \varepsilon.$$

uniform convergence $\implies$ pointwise convergence
 $\nLeftarrow$

Uniform Convergence

The sense in which **uniform convergence** is better behaved than **pointwise convergence** is that it does preserve at least some properties of the sequence of functions.

Which properties?

Poll

- Go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on Math 3A03
- Click on Take Class Poll
- Fill in poll **Function sequences: Uniform convergence**
- .

Uniform Convergence

Theorem (Continuity and Uniform Convergence)

*Suppose $\{f_n\}$ is a sequence of functions that **converges uniformly** on $[a, b]$ to f . If each f_n is continuous on $[a, b]$, then f is continuous on $[a, b]$.*

What should our proof strategy be?

Our goal is to show that the limit function f is continuous for all $x \in [a, b]$. So given $x \in [a, b]$, we must show that for any $\varepsilon > 0$ we can find a small enough neighbourhood of x , say $(x - \delta, x + \delta)$ for some small δ , such that $|f(x) - f(y)| < \varepsilon$ if $y \in (x - \delta, x + \delta)$, i.e., if $|x - y| < \delta$.

Somewhat we have to manage this using the facts that (i) each f_n is continuous and (ii) $f_n \rightarrow f$ uniformly.

The key is that (for any n) if x and y are close then $f_n(x)$ and $f_n(y)$ are close, and, if n is large enough, f_n is (uniformly) close to f throughout $[a, b]$, so continuity is “passed through” to the limit.

Let's make this precise...

Uniform Convergence

Proof: f_n continuous $\forall n$ and $f_n \rightarrow f$ uniformly $\implies f$ continuous.

Fix $x \in [a, b]$ and $\varepsilon > 0$. We must show $\exists \delta > 0$ such that if $y \in [a, b]$ and $|x - y| < \delta$ then $|f(x) - f(y)| < \varepsilon$.

Since $f_n \rightarrow f$ uniformly, $\exists N \in \mathbb{N}$ $\exists$ $|f_N(y) - f(y)| < \frac{\varepsilon}{3} \quad \forall y \in [a, b]$ (in particular, $x \in [a, b]$, so we have $|f_N(x) - f(x)| < \frac{\varepsilon}{3}$).

Fix such an integer N .

Since f_N is continuous, there is some $\delta > 0$ such that if $y \in [a, b]$ satisfies $|x - y| < \delta$, then $|f_N(x) - f_N(y)| < \frac{\varepsilon}{3}$. For such y , we then have

$$\begin{aligned} |f(x) - f(y)| &= |f(x) - f_N(x) + f_N(x) - f_N(y) + f_N(y) - f(y)| \\ &\leq |f(x) - f_N(x)| + |f_N(x) - f_N(y)| + |f_N(y) - f(y)| \\ &< \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon, \end{aligned}$$

as required. □

Uniform Convergence

Theorem (Integrability and Uniform Convergence)

Suppose $\{f_n\}$ is a sequence of functions that *converges uniformly* on $[a, b]$ to f . If each f_n is *integrable* on $[a, b]$, then f is *integrable* and

$$\int_a^b f = \lim_{n \rightarrow \infty} \int_a^b f_n.$$



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 20
Sequences and Series of Functions II
Tuesday 3 March 2026

Announcements

- I have posted Test 1 and my solutions on the [course web site](#).
- I have added another (simpler) [example](#) showing pointwise convergence is insufficient to preserve integrability.

Last time...

Convergence of sequences of functions:

- Pointwise convergence
- Uniform convergence
- Theorem about continuity and uniform convergence

Uniform Convergence

Theorem (Integrability and Uniform Convergence)

Suppose $\{f_n\}$ is a sequence of functions that *converges uniformly* on $[a, b]$ to f . If each f_n is *integrable* on $[a, b]$, then f is *integrable* and

$$\int_a^b f = \lim_{n \rightarrow \infty} \int_a^b f_n. \quad (*)$$

(TBB §9.5.2, p. 571ff)

Note: To prove $(*)$, we will need the fact that if f is integrable then so is $|f|$,

and $\left| \int_a^b f \right| \leq \int_a^b |f|$. This “triangle inequality” is an excellent exercise.

Uniform Convergence

Lemma (A useful estimate for upper minus lower sums)

Let g, h be bounded on $[a, b]$. If $\sup_{x \in [a, b]} |h(x) - g(x)| \leq \delta$, then for every partition P of $[a, b]$,

$$U(h, P) - L(h, P) \leq (U(g, P) - L(g, P)) + 2\delta(b - a).$$

Proof.

Fix a partition $P = \{a = t_0 < t_1 < \cdots < t_m = b\}$. For each subinterval $I_i = [t_{i-1}, t_i]$ we have

$$\sup_{x \in I_i} h(x) \leq \sup_{x \in I_i} (g(x) + \delta) = \left(\sup_{x \in I_i} g(x) \right) + \delta,$$

and similarly

$$\inf_{x \in I_i} h(x) \geq \inf_{x \in I_i} (g(x) - \delta) = \left(\inf_{x \in I_i} g(x) \right) - \delta.$$

Multiplying by $\Delta t_i = t_i - t_{i-1}$ and summing over i ,

$$U(h, P) \leq U(g, P) + \delta(b - a) \quad \text{and} \quad L(h, P) \geq L(g, P) - \delta(b - a).$$

Subtract: $U(h, P) - L(h, P) \leq U(g, P) - L(g, P) + 2\delta(b - a).$ □

Uniform Convergence

Proof that f is **integrable** (using the ε - P criterion).

First, note that each f_n is bounded (integrable $\Rightarrow$ bounded). Since $f_n \rightarrow f$ uniformly, the limit function f is bounded too, so $U(f, P)$ and $L(f, P)$ make sense. We must show: $\forall \varepsilon > 0, \exists$ a partition P $\vdash U(f, P) - L(f, P) < \varepsilon$.

Given $\varepsilon > 0$, choose $\delta = \frac{\varepsilon}{4(b-a)}$. Since $f_n \rightarrow f$ uniformly, $\exists N \in \mathbb{N}$ such that

$$\sup_{x \in [a, b]} |f(x) - f_N(x)| < \delta.$$

Because f_N is **integrable**, the ε - P criterion gives a partition P such that

$$U(f_N, P) - L(f_N, P) < \frac{\varepsilon}{2}.$$

Apply the estimate from the **lemma** (previous slide) with $g = f_N$ and $h = f$:

$$\begin{aligned} U(f, P) - L(f, P) &\leq (U(f_N, P) - L(f_N, P)) + 2\delta(b-a) \\ &< \frac{\varepsilon}{2} + 2 \cdot \frac{\varepsilon}{4(b-a)}(b-a) = \varepsilon. \end{aligned}$$

This is exactly the ε - P criterion for f to be **integrable**. □

Now that we know f is **integrable**, we can prove (*) ...

Uniform Convergence

Proof that $\int_a^b f = \lim_{n \rightarrow \infty} \int_a^b f_n$ given that f is integrable.

Given that f is **integrable**, to prove the equality, we will show that

$$\forall \varepsilon > 0, \quad \exists N \in \mathbb{N} \quad \text{such that} \quad \left| \int_a^b f - \int_a^b f_n \right| < \varepsilon \quad \forall n \geq N.$$

For any $n \in \mathbb{N}$, we have

$$\begin{aligned} \left| \int_a^b f - \int_a^b f_n \right| &= \left| \int_a^b (f - f_n) \right| \leq \int_a^b |f - f_n| \\ &\leq U(|f - f_n|, \{a, b\}) = \left(\sup_{x \in [a, b]} |f(x) - f_n(x)| \right) (b - a). \end{aligned}$$

But f_n **converges uniformly** to f , which means that


$$\exists N \in \mathbb{N} \quad \text{such that} \quad \sup_{x \in [a, b]} |f(x) - f_n(x)| < \frac{\varepsilon}{b - a} \quad \forall n \geq N.$$

For such n , we have $\left| \int_a^b f - \int_a^b f_n \right| < \varepsilon$, as required. □

Uniform Convergence

The interaction between **uniform convergence** and differentiability is more subtle.

Example (f_n diff'ble $\forall n$ and $f_n \rightarrow f$ uniformly $\not\Rightarrow f$ diff'ble)



$$f_n(x) = \frac{1}{2n} + (x^2)^{(1+\frac{1}{n})/2}$$

Each f_n is differentiable

$$f_n(x) \rightarrow f(x) = |x|$$

uniformly


Limit function f is not differentiable.

Note: Graph shows $n = 1, 2, 4, 64$ for $x \in [-1, 1]$.

Uniform Convergence

Even if $f_n \rightarrow f$ uniformly, and all the f_n and f are differentiable, it is not necessarily true that $f'_n \rightarrow f'$.

Example ($f_n \rightarrow f$ uniformly and f'_n, f' exist $\not\Rightarrow f'_n \rightarrow f'$)



$$f_n(x) = \frac{1}{n} \sin(n^2 x)$$

$$f_n(x) \rightarrow f(x) \equiv 0$$

uniformly

$$f'_n(x) = n \cos(n^2 x)$$

$$\lim_{n \rightarrow \infty} f'_n(x) \text{ does not exist}$$

(e.g., $f'_n(0) = n$, which diverges as $n \rightarrow \infty$)

Note: Graph shows $n = 1, 2$ on interval $[-20, 20]$.

Uniform Convergence

The [theorem on integrability and uniform convergence](#), together with the [fundamental theorem of calculus](#), must yield some result on uniformly convergent sequences of differentiable functions.

But the result must make hypotheses that avoid the failures in the examples on the two previous slides.

Theorem (Differentiability and Uniform Convergence)

Suppose $\{f_n\}$ is a sequence of differentiable functions on $[a, b]$ such that

- 1** f'_n is integrable for each n ,
- 2** the sequence $\{f'_n\}$ converges [uniformly](#) on $[a, b]$ to a continuous function g ,
- 3** the sequence $\{f_n\}$ converges [pointwise](#) to a function f .

Then f is differentiable and $\{f'_n\}$ converges [uniformly](#) to f' .



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 21
Sequences and Series of Functions III
Thursday 5 March 2026

Announcements

- Assignment 4 is posted on the course web site.

Last time: Convergence of sequences of functions:

- Pointwise convergence
- Uniform convergence
- Theorem about continuity and uniform convergence
- Theorem about integrability and uniform convergence
- Theorem about differentiability and uniform convergence

Uniform convergence

Proof of theorem on Differentiability and Uniform Convergence.

Since the function g to which f'_n converges is continuous, it is certainly integrable. So we can apply the [theorem on integrability and uniform convergence](#) on the interval $[a, x]$ to infer that

$$\begin{aligned}\int_a^x g &= \lim_{n \rightarrow \infty} \int_a^x f'_n && f'_n \rightarrow g \text{ uniformly} \\ &= \lim_{n \rightarrow \infty} (f_n(x) - f_n(a)) && \text{SFTC} \\ &= f(x) - f(a) && f_n \rightarrow f \text{ pointwise}\end{aligned}$$

Since g is continuous, [FFTC](#) implies that

$$g(x) = \lim_{n \rightarrow \infty} f'_n(x) = f'(x)$$

for all $x \in [a, b]$. □

Series of Functions

Series of Real Numbers

Suppose $\{x_n\}$ is a sequence of real numbers. Recall that the *sequence of partial sums* is the sequence $\{s_n\}$ defined by

$$s_n = \sum_{k=1}^n x_k .$$

If the sequence of partial sums converges, then we write the limit as

$$\sum_{k=1}^{\infty} x_k = \lim_{n \rightarrow \infty} \sum_{k=1}^n x_k = \lim_{n \rightarrow \infty} s_n .$$

In this case, we call $\sum_{k=1}^{\infty} x_k$ a *convergent series*. A *divergent series* is a sequence of partial sums that diverges; we sometimes abuse notation and write $\sum_{k=1}^{\infty} x_k$ for divergent series as well.

A *series* is either a convergent series or a divergent series.

Our goal now is to extend this notion of series to sequences of functions.

Series of Functions

Suppose $\{f_n\}$ is a sequence of functions defined on a set $D \subseteq \mathbb{R}$. The **sequence of partial sums** is the sequence $\{S_n\}$ where S_n is the function defined on D by

$$S_n(x) = \sum_{k=1}^n f_k(x).$$

When talking about limits of the S_n , we will write $\sum_{k=1}^{\infty} f_k$ and refer to this as a **series**.

Keep in mind that this is very informal, since the terminology does not specify any sense in which the S_n converge, nor does it assume that the S_n converge at all!

We will now make this more formal.

Series of Functions

Suppose $\{f_n\}$ is a sequence of functions defined on a domain D , and $\{S_n\}$ is its **sequence of partial sums**.

Definition (Convergence of Series)

If the sequence of partial sums $\{S_n\}$ **converges pointwise** on D to a function f , then we say that the series $\sum_{k=1}^{\infty} f_k$ **converges pointwise on D to f** .

If the $\{S_n\}$ **converge uniformly** on D to a function f , then we say that the series $\sum_{k=1}^{\infty} f_k$ **converges uniformly on D to f** .

In both cases, we will write $f = \sum_{k=1}^{\infty} f_k$ to denote that the **series converges to f** .

Poll

- Go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on Math 3A03
- Click on Take Class Poll
- Fill in poll **Function sequences: Uniform convergence of series**
- .

Series of Functions

The theorems on convergence of sequences of **integrable**, **continuous** and **differentiable** functions have several immediate implications for series of functions.

In the following, we assume that $\{f_n\}$ is a sequence of functions defined on an interval $[a, b]$.

Corollary (Integrals of Series)

*Suppose the f_n are **integrable** and $\sum_{k=1}^{\infty} f_k$ **converges uniformly** to a function f . Then f is **integrable** and*

$$\int_a^b f = \sum_{k=1}^{\infty} \int_a^b f_k.$$

Series of Functions

Corollary (Continuity of Series)

Suppose the f_n are continuous and $\sum_{k=1}^{\infty} f_k$ converges uniformly to a function f . Then f is continuous.

Corollary (Differentiability of Series)

Suppose $\{f_n\}$ is a sequence of differentiable functions on $[a, b]$ such that

- *f'_n is integrable for each n ,*
- *the series $\sum_{k=1}^{\infty} f'_k$ converges uniformly on $[a, b]$ to a continuous function g ,*
- *the series $\sum_{k=1}^{\infty} f_k$ converges pointwise to a function f .*

Then f is differentiable and $f' = \sum_{k=1}^{\infty} f'_k$.

Proving Uniform Convergence of Series of Functions

We have seen that several useful conclusions can be drawn when a series **converges uniformly**. The following gives a practical way of proving uniform convergence for a series of functions.

Theorem (Weierstrass M -test)

Let $\{f_n\}$ be a sequence of functions defined on $D \subseteq \mathbb{R}$, and suppose $\{M_n\}$ is a sequence of real numbers such that

$$|f_n(x)| \leq M_n, \quad \forall x \in D, \forall n \in \mathbb{N}.$$

*If $\sum_{n=1}^{\infty} M_n$ converges, then $\sum_{k=1}^{\infty} f_k$ converges **uniformly**.*

Proving Uniform Convergence of Series of Functions

Approach to proving the Weierstrass M-test:

- Let $S_n = \sum_{k=1}^n f_k$ be the n^{th} partial sum.
- Show that for every $\varepsilon > 0$, there is some $N \in \mathbb{N}$ so that

$$\sup_{x \in D} |S_n(x) - S_m(x)| < \varepsilon, \quad \forall n, m \geq N.$$

This condition is called the *uniform Cauchy criterion*.

- Prove that the uniform Cauchy criterion implies *uniform convergence*.
 - This part is an excellent exercise for you.

Note: The proof is similar to the proof of the *Cauchy criterion for real numbers*.

Proving Uniform Convergence of Series of Functions

Proof of the Weierstrass M -test.

Let $\varepsilon > 0$. Suppose the series $\sum M_n$ converges. By the **Cauchy criterion for real numbers**, there is some integer N so that

$$\left| \sum_{k=1}^n M_k - \sum_{k=1}^m M_k \right| < \varepsilon, \quad \forall n, m \geq N.$$

Without loss of generality, we can assume $m < n$, so the above can be written

$$M_{m+1} + M_{m+2} + \cdots + M_n < \varepsilon.$$

Note that we have $S_n - S_m = f_{m+1} + f_{m+2} + \cdots + f_n$, so the assumption that $|f_k(x)| \leq M_k$ implies that

$$\begin{aligned} |S_n(x) - S_m(x)| &= |f_{m+1}(x) + f_{m+2}(x) + \cdots + f_n(x)| \\ &\leq M_{m+1} + M_{m+2} + \cdots + M_n < \varepsilon. \end{aligned}$$

This is true $\forall x \in D$, hence $\sup_{x \in D} |S_n(x) - S_m(x)| < \varepsilon$, i.e.,

the uniform Cauchy criterion is satisfied. □

Proving Uniform Convergence of Series of Functions

In order to use the [Weierstrass M-test](#), we need to know whether an associated series of real numbers converges. The most useful standard results are:

- The geometric series $\sum_{n=0}^{\infty} a^n$ converges if and only if $|a| < 1$, in

which case the sum of the series is $\frac{1}{1-a}$.

- The **Ratio Test**: (TBB [Theorem 3.28](#))

If $a_n > 0$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} < 1$ then $\sum_{n=1}^{\infty} a_n$ converges.

- The **Root Test**: (TBB [Theorem 3.30](#))

If $a_n \geq 0$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} < 1$ then $\sum_{n=1}^{\infty} a_n$ converges.

Proving Uniform Convergence of Series of Functions

- The ***Integral Test***: (TBB Theorem 3.35)

Let f be a nonnegative decreasing function on $[1, \infty)$ such that

$$\int_1^K f \text{ exists for all } K > 1.$$

Then

$$\sum_{k=1}^{\infty} f(k) \text{ converges} \iff \lim_{K \rightarrow \infty} \int_1^K f(x) dx \text{ exists.}$$

(*There are many other known results concerning convergence series of real numbers.* See TBB Chapter 3.)

Proving Uniform Convergence of Series of Functions

Example

Let $p > 1$, and consider the series $\sum_{k=1}^{\infty} \frac{\sin(kx)}{k^p}$.

This series satisfies $\left| \frac{\sin(kx)}{k^p} \right| \leq \frac{1}{k^p}$ for all $x \in \mathbb{R}$.

Since the series $\sum_{k=1}^{\infty} \frac{1}{k^p}$ converges (by the integral test), it follows from the [Weierstrass M-test](#) that the series $\sum_{k=1}^{\infty} \frac{\sin(kx)}{k^p}$ [converges uniformly](#).
[Hence](#) it is a continuous function.

In fact, if $p > 2$ then the series $\sum_{k=1}^{\infty} \frac{\sin(kx)}{k^p}$ is [differentiable](#):

Let $f_k(x) = \frac{\sin(kx)}{k^p}$. The f'_k are continuous and another application of the [Weierstrass M-test](#) shows that $\sum_{k=1}^{\infty} f'_k$ converges uniformly. Hence the series is differentiable and the derivative is $\sum_{k=1}^{\infty} f'_k$.



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 22
Sequences and Series of Functions IV
Friday 6 March 2026