

24 Differentiation

25 Differentiation II

26 Differentiation II

Differentiation



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I


Instructor: David Earn

Lecture 24
Differentiation
Tuesday 5 November 2019

Announcements

- [Assignment 4](#) is posted and is due on Tuesday 12 Nov 2019, 2:25pm, via [crowdmark](#).

The Derivative



The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The **derivative** of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If $f'(x_0)$ is finite we say that f is **differentiable** at x_0 . If f is differentiable at every point of a set $E \subseteq I$, we say that f is differentiable on E . If E is all of I , we simply say that f is a **differentiable function**.

Note: “Differentiable” and “a derivative exists” always mean that the derivative is finite.

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \lim_{x \rightarrow 2} \frac{(x + 2)(x - 2)}{x - 2} = \lim_{x \rightarrow 2} x + 2 = 4$$

Note:

- In the first two limits, we must have $x \neq 2$.
- But in the third limit, we just plug in $x = 2$.
- Two things are equal, but in one $x \neq 2$ and in the other $x = 2$.
- Good illustration of why it is important to define the meaning of limits rigorously.


Poll

- Go to https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Lecture 24: Differentiable at 0**
- .

The Derivative

Example

Let f be defined in a neighbourhood I of 0, and suppose $|f(x)| \leq x^2$ for all $x \in I$. Is f necessarily differentiable at 0? e.g.,



The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| = \left| \frac{x^2 \sin \frac{1}{x^2}}{x} \right| = \left| x \sin \frac{1}{x^2} \right| \leq |x|$$

Therefore:

$$|f'(0)| = \left| \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} \right| = \lim_{x \rightarrow 0} \left| \frac{f(x) - f(0)}{x - 0} \right| \leq \lim_{x \rightarrow 0} |x| = 0.$$

$\therefore f$ is differentiable at 0 and $f'(0) = 0$. □

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The **right-hand derivative** of f at x_0 , denoted by $f'_+(x_0)$, is the limit

$$f'_+(x_0) = \lim_{x \rightarrow x_0^+} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this one-sided limit exists or is infinite.

Similarly, the **left-hand derivative** of f at x_0 , denoted by $f'_-(x_0)$, is the limit

$$f'_-(x_0) = \lim_{x \rightarrow x_0^-} \frac{f(x) - f(x_0)}{x - x_0}.$$


Note:

If $x_0 \in I^\circ$ then f is differentiable at x_0 iff $f'_+(x_0) = f'_-(x_0) \neq \pm\infty$.

The Derivative

Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$



- Same slope from left and right. Why isn't f differentiable???
- $\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^+} f'(x) = \lim_{x \rightarrow 0} f'(x) = 1.$
- $f'_-(0) = f'_+(0) = f'(0) = \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \infty.$

The Derivative

- Higher derivatives: we write
 - $f'' = (f')'$ if f' is differentiable;
 - $f^{(n+1)} = (f^{(n)})'$ if $f^{(n)}$ is differentiable.

- Other standard notation for derivatives:

$$\frac{df}{dx} = f'(x)$$

$$D = \frac{d}{dx}$$

$$D^n f(x) = \frac{d^n f}{dx^n} = f^{(n)}(x)$$

The Derivative

Theorem (Differentiable \implies continuous)

If f is defined in a neighbourhood I of x_0 and f is differentiable at x_0 then f is continuous at x_0 .

Proof.

Must show $\lim_{x \rightarrow x_0} f(x) = f(x_0)$, i.e., $\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = 0$.

$$\begin{aligned}\lim_{x \rightarrow x_0} (f(x) - f(x_0)) &= \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \times (x - x_0) \right) \\ &= \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \right) \times \lim_{x \rightarrow x_0} (x - x_0) \\ &= f'(x_0) \times 0 = 0,\end{aligned}$$

where we have used the theorem on the algebra of limits. \square

The Derivative

Theorem (Algebra of derivatives)

Suppose f and g are defined on an interval I and $x_0 \in I$. If f and g are differentiable at x_0 then $f + g$ and fg are differentiable at x_0 . If, in addition, $g(x_0) \neq 0$ then f/g is differentiable at x_0 . Under these conditions:

- 1** $(cf)'(x_0) = cf'(x_0)$ for all $c \in \mathbb{R}$;
- 2** $(f + g)'(x_0) = (f' + g')(x_0)$;
- 3** $(fg)'(x_0) = (f'g + fg')(x_0)$;
- 4** $\left(\frac{f}{g}\right)'(x_0) = \left(\frac{gf' - fg'}{g^2}\right)(x_0) \quad (g(x_0) \neq 0).$

(Textbook (TBB) [Theorem 7.7, p. 408](#))

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$ such that $f(U) \subseteq V$. If f is differentiable at x_0 and g is differentiable at $f(x_0)$ then the composite function $h = g \circ f$ is differentiable at x_0 and

$$h'(x_0) = (g \circ f)'(x_0) = g'(f(x_0))f'(x_0).$$

(Textbook (TBB) §7.3.2, p. 411)

TBB provide a very good motivating discussion of this proof, which is quite technical.


The Derivative

Theorem (Derivative at local extrema)

Let $f : (a, b) \rightarrow \mathbb{R}$. If x is a maximum or minimum point of f in (a, b) , and f is differentiable at x , then $f'(x) = 0$.

(Textbook (TBB) [Theorem 7.18, p. 424](#))

Note: f need not be differentiable or even continuous at other points.





Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 25
Differentiation II
Thursday 7 November 2019

Poll

- Go to https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Test #1 Result**
- .

Announcements

- [Assignment 4](#) is posted and is due on Tuesday 12 Nov 2019, 2:25pm, via [crowdmark](#).
- [Test 1](#) has been returned via [crowdmark](#). Carefully read the solutions, which are posted on the course web site.

Last time...

- Definition of the **derivative**.
- Proved **differentiable** \implies **continuous**.
- Discussed **algebra of derivatives** and **chain rule**.
- Pictorial argument that **derivative is zero at extrema**.
- Defined **one-sided derivatives**
 - **Example**

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence $f'(x) = 0 \forall x \in (a, b)$. \square


Theorem (Mean value theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) then there exists $x \in (a, b)$ such that

$$f'(x) = \frac{f(b) - f(a)}{b - a}.$$

The Mean Value Theorem

Idea for proof:



Proof.

Apply [Rolle's theorem](#) to

$$h(x) = f(x) - \left[f(a) + \left(\frac{f(b) - f(a)}{b - a} \right) (x - a) \right].$$

□

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(x_*).$$

But $x_2 - x_1 > 0$ and since $x_* \in I$, we know $f'(x_*) > 0$.

$\therefore f(x_2) - f(x_1) > 0$, *i.e.*, $f(x_1) < f(x_2)$. □

Poll

- Go to https://www.childsmath.ca/childs/forms/main_login.php
- Click on [Math 3A03](#)
- Click on [Take Class Poll](#)
- Fill in poll **Lecture 25: IVP and derivatives**
- .

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the *intermediate value property* on I .

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the **intermediate value property** on I then it is impossible that it is the derivative of any function on I .
- Darboux's theorem implies that a derivative cannot have jump or removable discontinuities. Any discontinuity of a derivative must be **essential**. Recall example of a **discontinuous function with IVP**.

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the **Extreme Value Theorem** implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the **theorem on the derivative at local extrema**, we must have $f'(x) = 0$.

Now suppose more generally that $f'(a) < K < f'(b)$. Let

$g(x) = f(x) - Kx$. Then g is differentiable on I and $g'(x) = f'(x) - K$ for all $x \in I$. In addition, $g'(a) = f'(a) - K < 0$ and

$g'(b) = f'(b) - K > 0$, so by the argument above, $\exists x \in (a, b)$ such that $g'(x) = 0$, i.e., $f'(x) - K = 0$, i.e., $f'(x) = K$.

The case $f'(a) > K > f'(b)$ is similar. □



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 25
Differentiation II
Thursday 7 November 2019

Intermediate value property for derivatives

Example $(f'(x) \neq 0 \forall x \in I \implies f \nearrow \text{ or } \searrow \text{ on } I)$

If f is differentiable on an interval I and $f'(x) \neq 0$ for all $x \in I$ then f is either increasing or decreasing on the entire interval I .

Proof:

Suppose $\exists a, b \in I$ such that $f'(a) < 0$ and $f'(b) > 0$.

Then, from [Darboux's theorem](#), $\exists c \in I$ such that $f'(c) = 0$. $\implies \Leftarrow$

\therefore Either " $\exists a \in I \nmid f'(a) < 0$ " is FALSE
or " $\exists b \in I \nmid f'(b) > 0$ " is FALSE.

\therefore Since we know $f'(x) \neq 0 \forall x \in I$, it must be that
either $f'(x) > 0 \forall x \in I$ or $f'(x) < 0 \forall x \in I$,
i.e., either f is increasing on I or decreasing on I . □

The Derivative of an Inverse

Example (Sufficient condition for *non*-differentiable inverse)

Suppose f is continuous and one-to-one on an interval I . If $x \in I$ and $f'(x) = 0$ then f^{-1} is not differentiable at $y = f(x)$.

Proof: By definition, the inverse function satisfies

$$f(f^{-1}(y)) = y.$$

Suppose that f^{-1} is differentiable at y . Then, by the **Chain Rule**,

$$f'(f^{-1}(y)) \cdot (f^{-1})'(y) = 1.$$

But $f^{-1}(y) = x$, and $f'(x) = 0$, so

$$0 \cdot (f^{-1})'(y) = 1,$$

which is impossible! $\Rightarrow \Leftarrow$.



The Derivative of an Inverse

Theorem (Inverse function theorem)

If f is differentiable on an interval I and $f'(x) \neq 0 \forall x \in I$, then

1 f is one-to-one on I ;

2 f^{-1} is differentiable on $J = f(I)$;

3 $(f^{-1})'(f(x)) = \frac{1}{f'(x)}$ for all $x \in I$,

i.e., $(f^{-1})'(y) = \frac{1}{f'((f^{-1}(y)))}$ for all $y \in J$.

(Textbook (TBB) [Theorem 7.32](#), p. 445)

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

■ Darboux's theorem $\implies f \nearrow$ or \searrow on $I \implies f$ is 1 : 1 on I

■ If $y = f(x)$ and $y_0 = f(x_0)$

then $x = f^{-1}(y)$ and $x_0 = f^{-1}(y_0)$,

$$\begin{aligned} \text{so } \frac{f^{-1}(y) - f^{-1}(y_0)}{y - y_0} &= \frac{x - x_0}{f(x) - f(x_0)} \\ &= \frac{1}{\frac{f(x) - f(x_0)}{x - x_0}}. \end{aligned}$$

■ Since f continuous at x_0 , we know $x \rightarrow x_0 \implies y \rightarrow y_0$.

■ But we need $y \rightarrow y_0 \implies x \rightarrow x_0$, i.e., f^{-1} continuous at y_0 .

■ In fact, f continuous and either \nearrow or \searrow on $I \implies f^{-1}$ continuous on $J = f(I)$. (more generally, cf. [Invariance of Domain](#) thm)

Please consider. . .

5 minute *Student Respiratory Illness Survey:*

<https://surveys.mcmaster.ca/limesurvey/index.php/893454>

Please complete this anonymous survey to help us monitor the patterns of respiratory illness, over-the-counter drug use, and social contact within the McMaster community. There are no risks to filling out this survey, and your participation is voluntary. You do not need to answer any questions that make you uncomfortable, and all information provided will be kept strictly confidential. Thanks for participating.

–Dr. Marek Smieja (Infectious Diseases)