

- 22 Metric Spaces
- 23 Metric Spaces II
- 24 Metric Spaces III
- 25 Metric Spaces IV
- 26 Metric Spaces V
- 27 Metric Spaces VI
- 28 Metric Spaces VII



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 22
Metric Spaces
Monday 10 March 2025

Announcements

- New, exciting topic today...

Metric Spaces

The metric structure of \mathbb{R}

Definition (Absolute Value function)

For any $x \in \mathbb{R}$,

$$|x| \stackrel{\text{def}}{=} \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

Theorem (Properties of the Absolute Value function)

For all $x, y \in \mathbb{R}$:

- 1 $-|x| \leq x \leq |x|.$
- 2 $|xy| = |x| |y|.$
- 3 $|x + y| \leq |x| + |y|.$
- 4 $|x| - |y| \leq |x - y|.$

The metric structure of \mathbb{R}

Definition (Distance function or metric)

The distance between two real numbers x and y is

$$d(x, y) = |x - y| .$$

Theorem (Properties of distance function or metric)

- $d(x, y) \geq 0$ *distances are positive or zero*
- $d(x, y) = 0 \iff x = y$ *distinct points have distance > 0*
- $d(x, y) = d(y, x)$ *distance is symmetric*
- $d(x, y) \leq d(x, z) + d(z, y)$ *the triangle inequality*

Note: Any function satisfying these properties can be considered a “distance” or “metric”.

The metric structure of \mathbb{R}

Given $d(x, y) = |x - y|$, the **properties of the distance function** are equivalent to:

Theorem (Metric properties of the absolute value function)

For all $x, y \in \mathbb{R}$:

- 1 $|x| \geq 0$
- 2 $|x| = 0 \iff x = 0$
- 3 $|x| = |-x|$
- 4 $|x + y| \leq |x| + |y|$ (*the triangle inequality*)

Slick proof of the triangle inequality

Theorem (The Triangle Inequality for the standard metric on \mathbb{R})

$|x + y| \leq |x| + |y|$ for all $x, y \in \mathbb{R}$.

Proof.

Let $s = \text{sign}(x + y)$. Then

$$|x + y| = s(x + y) = sx + sy \leq |x| + |y| ,$$

as required. □

A non-standard metric on \mathbb{R}

Example (finite distance between every pair of real numbers)

Let $f(x) = \frac{x}{1+x}$, and define $d(x, y) = f(|x - y|)$. Prove that $d(x, y)$ can be interpreted as a distance between x and y because it satisfies **all the properties of a metric**.

Proof: The only metric property that is non-trivial to prove is the triangle inequality. Note that $f(x)$ is an increasing function on $[0, \infty)$, so the usual triangle inequality, $|x - y| \leq |x - z| + |z - y|$, implies

$$\begin{aligned} f(|x - y|) &\leq f(|x - z| + |z - y|) = \frac{|x - z| + |z - y|}{1 + |x - z| + |z - y|} \\ &= \frac{|x - z|}{1 + |x - z| + |z - y|} + \frac{|z - y|}{1 + |x - z| + |z - y|} \\ &\leq \frac{|x - z|}{1 + |x - z|} + \frac{|z - y|}{1 + |z - y|} = f(|x - z|) + f(|z - y|) \end{aligned}$$

i.e., $d(x, y) \leq d(x, z) + d(z, y)$. □

Poll

- Go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Metric spaces: Is “= vs \neq ” a metric?**
- .

Discrete metric

Example (Discrete metric on \mathbb{R})

Let $d(x, y) = \begin{cases} 0 & x = y, \\ 1 & x \neq y. \end{cases}$ Is d a **metric** on \mathbb{R} ?

By definition, $d(x, y)$ is non-negative, zero iff $x = y$, and symmetric. For the triangle inequality, if $x = y$ then $d(x, y) = 0$ so the inequality holds for any z . If $x \neq y$ then $d(x, y) = 1$, and at least one of x and y must not equal z , so the inequality says either $1 \leq 1$ or $1 \leq 2$.

Example (Discrete metric on any set X)

The argument that $d(x, y)$ is a metric on \mathbb{R} has nothing to do with \mathbb{R} specifically. $d(x, y)$ is a metric on any set X .

General metric space (X, d)

Definition (Metric space)

A **metric space** (X, d) is a non-empty set X together with a distance function (or **metric**) $d : X \times X \rightarrow \mathbb{R}$ satisfying

- $d(x, y) \geq 0$ *distances are positive or zero*
- $d(x, y) = 0 \iff x = y$ *distinct points have distance > 0*
- $d(x, y) = d(y, x)$ *distance is symmetric*
- $d(x, y) \leq d(x, z) + d(z, y)$ *the triangle inequality*

Much of our analysis of sequences of real numbers and topology of \mathbb{R} generalizes to any metric space. Very often, definitions and proofs depend only on the the existence of a metric, not on $|x|$ specifically. Many useful inferences can be made by identifying a metric on a space of interest.

Examples of metric spaces

Example (Metric spaces (X, d))

- $X = \mathbb{Q}$, with the standard metric $d(x, y) = |x - y|$.
As $\mathbb{Q} \subset \mathbb{R}$, each **condition for d** is satisfied in \mathbb{Q} .

How different is (\mathbb{Q}, d) from (\mathbb{R}, d) ?

- $X = \mathbb{N}$, with the standard metric $d(x, y) = |x - y|$.
As $\mathbb{N} \subset \mathbb{R}$, each **condition for d** is satisfied in \mathbb{N} .
- $X = \mathbb{R}^2$ with $d(x, y) = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}$, where we write the vectors $x = (x_1, x_2)$, $y = (y_1, y_2) \in \mathbb{R}^2$.
- $X = \mathbb{R}^n$ with $d(x, y) = \sqrt{\sum_{i=1}^n (x_i - y_i)^2}$, where $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n) \in \mathbb{R}^n$.

This metric on \mathbb{R}^n is called the ***Euclidean distance***.

Metrics from norms

The Euclidean metric on \mathbb{R}^n is the (Euclidean) length of the difference of two vectors. This connection between length and distance generalizes to any vector space in which *length* is defined.

Definition (Norm)

A **norm** on a vector space X is a real-valued function on X such that if $x, y \in X$ and $\alpha \in \mathbb{R}$ then

- 1 $\|x\| \geq 0$ and $\|x\| = 0$ iff x is the zero element in X ;
- 2 $\|\alpha x\| = |\alpha| \|x\|$;
- 3 $\|x + y\| \leq \|x\| + \|y\|$.

A vector space X equipped with a norm $\|\cdot\|$ is said to be a **normed vector space**. Any norm $\|\cdot\|$ **induces** a metric d via

$$d(x, y) = \|x - y\|.$$

Proving that a function is a norm is not necessarily easy. Let's try for the Euclidean norm... To that end, recall the notion of inner product...

Norms from inner products

Definition (Inner product)

An *inner product* on a vector space V over \mathbb{R} is a function

$$\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$$

such that for all $u, v, w \in V$ and all scalars $\alpha \in \mathbb{R}$:

- $\langle u, v \rangle = \overline{\langle v, u \rangle}$ *conjugate symmetry*
- $\langle \alpha u + v, w \rangle = \alpha \langle u, w \rangle + \langle v, w \rangle$ *linearity in 1st argument*
- $\langle v, v \rangle \geq 0$ with equality iff $v = 0$ *positive definiteness*

A vector space equipped with an inner product is called an *inner product space*.

Definition (Inner Product Norm)

The norm induced by an inner product $\langle \cdot, \cdot \rangle$ is $\|u\| = \sqrt{\langle u, u \rangle}$.

Norms from inner products

Theorem (Cauchy-Schwarz inequality)

Let V be a (real) inner product space with inner product $\langle \cdot, \cdot \rangle$. For all vectors $u, v \in V$, we have

$$|\langle u, v \rangle| \leq \|u\| \|v\|$$

where $\|u\| = \sqrt{\langle u, u \rangle}$ is the norm induced by the inner product.

Proof.

The standard proof begins with an idea that probably took someone a long time to think of: Since $\langle v, v \rangle \geq 0$ for any $v \in V$, for any $t \in \mathbb{R}$ we have

$$\begin{aligned} 0 &\leq \langle u + tv, u + tv \rangle = \langle u, u \rangle + t \langle u, v \rangle + t \langle v, u \rangle + t^2 \langle v, v \rangle \\ &= \langle u, u \rangle + 2t \langle u, v \rangle + t^2 \langle v, v \rangle \end{aligned}$$

This is a quadratic polynomial in t , which is non-negative for all $t \in \mathbb{R}$. Hence, this quadratic has at most one real root. Consequently, its discriminant is non-positive, *i.e.*, $(2 \langle u, v \rangle)^2 - 4 \langle u, u \rangle \langle v, v \rangle \leq 0$.

continued...

Norms from inner products

Proof of Cauchy-Schwarz inequality (continued).

Simplifying the non-positive discriminant condition, we have

$$(\langle u, v \rangle)^2 \leq \langle u, u \rangle \langle v, v \rangle .$$

Taking square roots, we have

$$|\langle u, v \rangle| \leq \|u\| \|v\| ,$$

as required. □

How might you come up with such a proof?

Perhaps by guessing the result (based on knowing it in \mathbb{R}^2) and then working backwards.



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 23
Metric Spaces II
Wednesday 12 March 2025

Announcements

- Participation deadline for Assignment 4 was at 11:25am today.

Last time...

- Introduction to **metric spaces**
 - Critical ingredient: the triangle inequality
- **Cauchy-Schwarz inequality**
(proved for real inner product spaces)

Norms from inner products

If X is an inner product space, then Cauchy-Schwarz allows us to prove that the induced norm really is a norm (i.e., satisfies the triangle inequality). For $x, y \in X$, we have

$$\begin{aligned}\|x + y\|^2 &= \langle x + y, x + y \rangle \\ &= \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle \\ &= \|x\|^2 + \|y\|^2 + 2 \langle x, y \rangle \\ &\leq \|x\|^2 + \|y\|^2 + 2 \|x\| \|y\| \\ &= (\|x\| + \|y\|)^2\end{aligned}$$

$$\implies \|x + y\| \leq \|x\| + \|y\| .$$

In particular, \mathbb{R}^n with the usual “dot product” $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$ induces the Euclidean norm $\|\cdot\|_2$, which therefore really is a norm, and $d(x, y) = \|x - y\|_2$ really is a metric (the Euclidean distance).

What about other norms induced by inner products?

Other metric spaces induced by inner products

We are accustomed to finite vectors:

$$\begin{aligned}x &= (x_1, x_2) \in X = \mathbb{R}^2 \\x &= (x_1, x_2, x_3) \in X = \mathbb{R}^3 \\x &= (x_1, x_2, \dots, x_n) \in X = \mathbb{R}^n\end{aligned}$$

We can think of a sequence as an infinite vector:

$$x = (x_1, x_2, \dots) \in X = \{\{x_n\} : n \in \mathbb{N}\}$$

The points in this space (X) are infinite-dimensional vectors.

We can think of an infinite vector as a function:

$$x_n = f(n) \quad \implies \quad (x_1, x_2, \dots) = (f(1), f(2), \dots)$$

The points in this space are functions: $X = \{f \mid f : \mathbb{N} \rightarrow \mathbb{R}\}$.

So we can generalize to other spaces via functions, e.g.,

$$C[0, 1] = \{f : [0, 1] \rightarrow \mathbb{R} \mid f \text{ continuous}\}$$

All of the above spaces have a natural inner product, and hence a natural norm and metric.

Other metric spaces induced by inner products

Inner products that convert the spaces on the previous slide into (Euclidean) metric spaces:

$$\mathbb{R}^n: \quad \langle x, y \rangle = \sum_{i=1}^n x_i y_i$$

$$\ell^2(\mathbb{R}): \quad \langle x, y \rangle = \sum_{i=1}^{\infty} x_i y_i$$

$$C[a, b]: \quad \langle f, g \rangle = \int_a^b f(x)g(x) dx$$

Note: ℓ^2 includes only **square-summable** sequences: $\sum_{n=1}^{\infty} x_n^2 < \infty$


*Do we need to specify that $C[a, b]$ contains only **square-integrable** functions?*

Metrics from norms

Example (Taxicab distance)

Taxicab norm on \mathbb{R}^n

$$\|x\|_1 = \sum_{i=1}^n |x_i|$$



In taxicab geometry, the lengths of the **red**, **blue**, **green**, and **yellow** paths all equal 12, the taxicab distance between the opposite corners, and all four paths are shortest paths. Instead, in Euclidean geometry, the **red**, **blue**, and **yellow** paths still have length 12 but the **green** path is the unique shortest path, with length equal to the Euclidean distance between the opposite corners, $6\sqrt{2} \approx 8.49$.

Image and caption from [Wikipedia article on "Taxicab geometry"](#).

Note: The **green** path can be followed. All the points of \mathbb{R}^2 are still present when we measure distance with the taxicab metric. Any monotonic curve path that joins the two points can be approximated by an arbitrarily fine grid, and will have the same length. The metric does not impose a particular grid.

Metrics from norms

Example (p -metric)

$$p\text{-norm on } \mathbb{R}^n \text{ (for } p \geq 1) \quad \|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}.$$

- $p = 1$ is the taxicab norm.
- $p = 2$ is the Euclidean norm.

What happens as $p \rightarrow \infty$? For any $x \in \mathbb{R}^n$, we have

$$\begin{aligned} \|x\|_p &= \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} = |x_k| \left(\sum_{i=1}^n \left| \frac{x_i}{x_k} \right|^p \right)^{\frac{1}{p}} \quad (|x_k| > |x_i| \quad \forall i \neq k) \\ &= |x_k| \left(1 + \sum_{i \neq k} \left| \frac{x_i}{x_k} \right|^p \right)^{\frac{1}{p}} \xrightarrow{p \rightarrow \infty} |x_k| \end{aligned}$$

What further work is required if $\nexists k \} |x_k| > |x_i| \quad \forall i \neq k$?

Therefore, we define $\|\cdot\|_\infty$ to be

$$\text{Max norm on } \mathbb{R}^n \quad \|x\|_\infty = \max_{1 \leq i \leq n} |x_i|$$

Poll

- Go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Metric spaces: Which p -norms are induced?**
- .

Metrics from norms

Example (p -metric)

Proving that $p = 1$ and $p = \infty$ yield norms is a *good exercise*.

Only $p = 2$ is induced by an inner product.

That the p -norms for $p \neq 1, 2, \infty$ are norms is harder to prove (but true), so

$$d_p(x, y) = \|x - y\|_p$$

is a metric on \mathbb{R}^n for any $p \geq 1$.

Topology of metric spaces

We can generalize the notion of “neighbourhood of a point” to any metric space:

Definition (Open ball)

Let (X, d) be a metric space. If $x_0 \in X$ and $r > 0$ then the **open ball of radius r about x_0** is

$$B_r(x_0) = \{x \in X : d(x, x_0) < r\}.$$

x_0 is said to be the **centre** of $B_r(x_0)$.

Note: The notation $B(x_0, r)$ is also common (and used in TBB).


Definition (Neighbourhood)

A **neighborhood** of x is any set that contains an open ball $B_r(x)$ for some $r > 0$.

Topology of metric spaces

Example (Open balls in metric spaces)

- In the metric space $(\mathbb{R}, \text{standard})$, i.e., \mathbb{R} with $d(x, y) = |x - y|$, $B_r(x) = (x - r, x + r)$, an open interval of length $2r$ centred at x .
- In \mathbb{R}^n with Euclidean metric $d(x, y) = \|x - y\|_2$, $B_r(x)$ has a spherical boundary (circular boundary if $n = 2$).
- In \mathbb{R}^n with a p -norm $\|\cdot\|_p$, the ball is not spherical. For \mathbb{R}^2 with the Taxicab metric $d(x, y) = \|x - y\|_1$, $B_r(x)$ is diamond shaped, and for \mathbb{R}^2 with the Max norm $\|\cdot\|_\infty$, $B_r(x)$ is a square. We write $B_r^p(x)$ for balls in the p -norm (" p -balls").



$$B_r^p(x)$$

- $p = 1$
- $p = 2$
- $p = 16$

Topology of metric spaces

Example (p -norm inequalities and containments)

The following inequalities relate the various p -norms on \mathbb{R}^n ,

$$\|x\|_\infty \leq \|x\|_2 \leq \|x\|_1 \leq n \|x\|_\infty, \quad \text{and} \quad \|x\|_1 \leq \sqrt{n} \|x\|_2.$$

A good exercise.

Balls in the norm $\|\cdot\|_p$ are often written $B_r^p(x)$. The inequalities above imply that the following sets are *nested*:

$$B_{r/n}^2(x) \subset B_{r/n}^\infty(x) \subset B_r^1(x) \subset B_r^2(x) \subset B_r^\infty(x).$$

Another good exercise.

Example (Balls in the discrete metric)

For any set X , in the *discrete metric* the balls are simple, but strange. If $0 < r \leq 1$ then $B_r(x) = \{x\}$, a single point! If $r > 1$ then $B_r(x) = X$, the whole space! You can't be "close" to a point x unless you are at x itself!



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 25
Metric Spaces III
Monday 17 March 2025

Announcements

Discussion of metrics so far. . .

- Metrics induced by norms, e.g., p -norms
- Metrics from norms induced by inner products
- Metric on any set: discrete metric
- Balls (p -balls, discrete balls)

Note:

- I added a note to the slide on the taxicab metric.

Topology of metric spaces

Definition (Convergence of a sequence in a metric space)

Let (X, d) be any **metric space**. A sequence $(x_n)_{n \in \mathbb{N}}$ **converges** to $x \in X$, written $x_n \xrightarrow{n \rightarrow \infty} x$ or $\lim_{n \rightarrow \infty} x_n = x$, if:

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \quad \} \quad d(x_n, x) < \varepsilon \quad \forall n \geq N.$$

If the sequence does not converge to any $x \in X$, we say it **diverges**.

Equivalently, $x_n \xrightarrow{n \rightarrow \infty} x$ if, for any ball $B_\varepsilon(x)$ centered at x , the sequence (x_n) lies inside that ball eventually ($\exists N \in \mathbb{N}$ such that $x_N \in B_\varepsilon(x)$), and stays inside it ($x_n \in B_\varepsilon(x) \forall n > N$), i.e.,

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \quad \} \quad x_n \in B_\varepsilon(x) \quad \forall n \geq N.$$

Topology of metric spaces

Definition (Boundedness in a metric space)

In any **metric space** (X, d) , a sequence (x_n) is **bounded** if there exists $x_0 \in X$ and $r > 0$ such that $x_n \in B_r(x_0)$ for all $n \in \mathbb{N}$.

Theorem

*In any **metric space** (X, d) , any convergent sequence is bounded.*

Note: The converse is FALSE. e.g., $x_n = (-1)^n$ in $(\mathbb{R}, \text{standard})$.

Proof.

Taking $\varepsilon = 1$ (or any particular value) in the definition of convergence, since $x_n \xrightarrow{n \rightarrow \infty} x$ in (X, d) , $\exists N \in \mathbb{N}$ with $d(x_n, x) < \varepsilon = 1$, $\forall n \geq N$, i.e., $x_n \in B_1(x)$, $\forall n \geq N$. The earlier elements of the sequence, x_1, \dots, x_{N-1} , are a finite collection, so we can choose $r > 0$ so that

$$r > \max\{d(x_1, x), d(x_2, x), \dots, d(x_{N-1}, x), 1\}.$$

With this r , $x_n \in B_r(x)$ holds $\forall n \in \mathbb{N}$. So (x_n) is bounded. \square

Topology of metric spaces

Definition (Interior point)

$x \in E \subseteq X$ is an **interior point** of the set E in the metric space (X, d) if x lies in an **open ball** that is contained in E , i.e.,

$$\exists \varepsilon > 0 \quad \text{) } B_\varepsilon(x) \subset E.$$

Definition (Interior of a set)

If $E \subseteq X$ then the **interior** of E , denoted $\text{int}(E)$ or E° , is the set of all interior points of E .

Note:

- $x \in E^\circ$ means not only that $x \in E$, but that there is an entire open ball $B_\varepsilon(x) \subset E$.
- For any *smaller* radius, $0 < r < \varepsilon$, we have $x \in B_r(x) \subseteq B_\varepsilon(x) \subseteq E$, so the choice of ε is not unique for an interior point.

Topology of metric spaces

Definition (Open set)

A set $E \subseteq X$ is **open** if every point of E is an **interior point**.

Example (Upper half-plane in $(\mathbb{R}^2, \text{Euclidean})$)

Let $W = \{x = (x_1, x_2) : x_2 > 0\}$. Then $\forall x \in W$, if $0 < \varepsilon < x_2$ then $B_\varepsilon(x) \subset W$. Thus all $x \in W$ are interior points. So W is an open set.

Example (Any set in $(X, \text{discrete})$)

$(X, \text{discrete})$ means X is a non-empty set and d is the discrete metric d . So, $\forall x \in X$,

$$B_\varepsilon(x) = \begin{cases} \{x\}, & \text{if } 0 < \varepsilon \leq 1, \\ X, & \text{if } \varepsilon > 1. \end{cases}$$

Therefore, for any subset $E \subseteq X$ and for any point $x \in E$, $B_1(x) = \{x\} \subseteq E$, so every point of any set E is an interior point, so **all** sets in the discrete metric are open sets!

Topology of metric spaces

Definition (Closed set)

A set $F \subseteq X$ is **closed** if F^c is **open**.

Example (Lower half-plane in $(\mathbb{R}^2, \text{Euclidean})$)

Let $Z = \{x = (x_1, x_2) : x_2 \leq 0\}$. Then $Z = W^c$, where W is the (**open**) upper half-plane. So Z is a closed set.

Note: In $(\mathbb{R}^2, \text{Euclidean})$, a half-plane is open or closed depending on whether none or all of the x -axis is included. If neither none nor all of the x -axis is included then the half-plane is neither open nor closed.

Example (Any set in $(X, \text{discrete})$)

Any set $E \subseteq X$ is **open**. So given any $F \subseteq X$, $F^c \subseteq X$ so F^c is open. Hence F is closed. So **all** sets are **both open and closed** with respect to the discrete metric.

Topology of metric spaces

When studying $X = \mathbb{R}$, we defined closed sets differently.

Could we have used the same definition for a general metric space?

Definition (Accumulation Point or Limit Point or Cluster Point)

If $E \subseteq X$ in a metric space (X, d) then x is an **accumulation point** of E if every **neighbourhood** of x contains infinitely many points of E ,

$$\text{i.e.,} \quad \forall \varepsilon > 0 \quad B_\varepsilon(x) \cap (E \setminus \{x\}) \neq \emptyset.$$

Equivalently, $x \in X$ is an accumulation point of the set E if and only if there exists a sequence $(x_n)_{n \in \mathbb{N}}$ with $x_n \in E \forall n \in \mathbb{N}$, such that $x_n \neq x \forall n$ and $x_n \xrightarrow{n \rightarrow \infty} x$.

Topology of metric spaces

Theorem (Closed sets in a metric space)

A set $F \subseteq X$ in a *metric space* (X, d) is *closed* if and only if F contains all its *accumulation points*.

Proof.

First, suppose F is closed, and that $x \in X$ is an accumulation point of F . If $x \notin F$, then $x \in F^c$, which is open. Therefore, $\exists \varepsilon > 0$ such that $B_\varepsilon(x) \subset F^c$, i.e., $B_\varepsilon(x) \cap F = \emptyset$. But this contradicts x being an accumulation point of F . So we must have $x \in F$.

Conversely, suppose F contains all its accumulation points. Then, if $z \in F^c$, z cannot be an accumulation point of F . But for any $\varepsilon > 0$, $z \in B_\varepsilon(z)$, so $\exists \varepsilon > 0$ for which $B_\varepsilon(z) \cap F = \emptyset$, i.e., $B_\varepsilon(z) \subset F^c$. Hence F^c is open, so F is closed. \square

Topology of metric spaces

Theorem (Properties of open sets in a metric space (X, d))

- 1 The sets X and \emptyset are open.
- 2 Any *intersection* of a *finite* number of open sets is open.
- 3 Any *union* of an *arbitrary* collection of open sets is open.
- 4 The complement of an open set is closed.

Theorem (Properties of closed sets in a metric space (X, d))

- 1 The sets X and \emptyset are closed.
- 2 Any *union* of a *finite* number of closed sets is closed.
- 3 Any *intersection* of an *arbitrary* collection of closed sets is closed.
- 4 The complement of a closed set is open.

Poll

- Go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Metric spaces: Set differences of balls**
- .

Topology of metric spaces

Example (Set differences of balls)

Suppose $0 < r_1 < r_2$ and (X, d) is a metric space. If $x \in X$, is it necessarily true that $B_{r_2}(x) \setminus B_{r_1}(x)$ is open or closed?

No, it depends on the metric d . If $(X, d) = (\mathbb{R}^n, \text{Euclidean})$ then $B_{r_2}(x) \setminus B_{r_1}(x)$ is neither open nor closed, e.g., in \mathbb{R}^1 ,

$$\begin{aligned} B_{r_2}(x) \setminus B_{r_1}(x) &= (x - r_2, x + r_2) \setminus (x - r_1, x + r_1) \\ &= (x - r_2, x - r_1] \cup [x + r_1, x + r_2). \end{aligned}$$

In contrast, if $(X, d) = (\mathbb{R}^n, \text{discrete})$ then $B_{r_2}(x) \setminus B_{r_1}(x)$ is open since any set in (X, d) is both open and closed.

In general, if A and B are sets then

$$A \setminus B = A \cap B^c,$$

so if A and B are both open, and B is also closed, then $A \setminus B$ is open.

Topology of metric spaces

Definition (Isolated point)

If $x \in E \subseteq X$ in a metric space (X, d) then x is an **isolated point** of E if there is a **neighbourhood** of x for which the only point in E is x itself, *i.e.*,

$$\exists \varepsilon > 0 \quad \vdash \quad B_\varepsilon(x) \cap E = \{x\}.$$

Example

Consider the metric space $(X, d) = ([0, 1], \text{standard})$.
What are the isolated points of (X, d) ?

There are no isolated points!

Now suppose $(X, d) = ([0, 1], \text{discrete})$.
What are the isolated points of (X, d) ?

All points are isolated!



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 26
Metric Spaces IV
Wednesday 19 March 2025

Announcements

Last time...

- More definitions and examples related to metric topology

Topology of metric spaces

Definition (Closure of a set)

If (X, d) is a metric space and $E \subseteq X$ then the **closure** of E , denoted \bar{E} , is the smallest closed set that contains E .

$\therefore \bar{E}$ is closed, $E \subseteq \bar{E}$, and if F is closed and $E \subseteq F$, then $\bar{E} \subseteq F$.

Theorem

$x \in \bar{E} \iff x$ is either an element of E or an **accumulation point** of E .
i.e., $x \in \bar{E} \iff \forall \varepsilon > 0, B_\varepsilon(x) \cap E \neq \emptyset$.

Proof.

(\Leftarrow) Suppose $\forall \varepsilon > 0, B_\varepsilon(x) \cap E \neq \emptyset$. In order to derive a contradiction, assume $x \notin \bar{E}$. Then $x \in (\bar{E})^c$, which is open, so $\exists \varepsilon > 0$ such that $B_\varepsilon(x) \subseteq (\bar{E})^c$, i.e., $B_\varepsilon(x) \cap \bar{E} = \emptyset$. But $E \subset \bar{E}$, so $B_\varepsilon(x) \cap E = \emptyset$. $\Rightarrow \Leftarrow$

(\Rightarrow) Suppose $x \in \bar{E}$ and, in order to derive a contradiction, suppose $\exists \varepsilon > 0$ $\} B_\varepsilon(x) \cap E = \emptyset$. Then $E \subseteq (B_\varepsilon(x))^c$. Hence $\bar{E} \subseteq \overline{(B_\varepsilon(x))^c} = (B_\varepsilon(x))^c$. But $x \in \bar{E}$ and $x \notin (B_\varepsilon(x))^c$. $\Rightarrow \Leftarrow$. $\therefore \forall \varepsilon > 0, B_\varepsilon(x) \cap E \neq \emptyset$. \square

Topology of metric spaces

Example (Cubic balls in \mathbb{R}^n)

In \mathbb{R}^n with the **max norm** $\|\cdot\|_\infty$, the distance between $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ is

$$d(x, y) = \|x - y\|_\infty = \max\{|x_i - y_i| : 1 \leq i \leq n\}.$$

Consider the set $E \subset \mathbb{R}^n$, where

$$E = \{x \in \mathbb{R}^n : 0 < x_i < 1, \forall i \in \{1, \dots, n\}\}.$$

What are the interior E° and closure \bar{E} of the set E ?


- **$n = 1$:** $E = (0, 1)$, an open interval. $E^\circ = (0, 1) = E$. $\bar{E} = [0, 1]$.
- **$n = 2$:** $E = (0, 1) \times (0, 1)$, an open square (an open ball in this metric).
 $E^\circ = (0, 1)^2 = E$. $\bar{E} = [0, 1]^2$.
- **$n > 2$:** $E = (0, 1)^n$, an open n -cube (an open ball in this metric).
 $E^\circ = (0, 1)^n = E$. $\bar{E} = [0, 1]^n$.

When we imagine an n -cube for $n > 3$, we are probably thinking about $n = 3$ in our minds. But we can easily represent individual points in \mathbb{R}^n in the plane. *How?*

Topology of metric spaces

Example (Cubic balls in \mathbb{R}^n)

Suppose $x, y \in E \subset \mathbb{R}^{28}$, $d(x, y) = \|x - y\|_\infty$.



The vertical **black line** indicates the distance between x and y .

Note: The same picture works for $\ell^\infty(\mathbb{R})$, the space of sequences that are bounded, and in which the norm is defined by sup rather than max.

Topology of metric spaces

For the space of continuous functions on a closed interval $[a, b]$, we defined the **Euclidean norm** via the standard inner product. We can also define a p -norm on this space for any $p \geq 1$,

$$\|f\|_p = \left(\int_a^b |f(x)|^p dx \right)^{1/p}.$$

As in finite dimensions, $\lim_{p \rightarrow \infty} \|f\|_p = \|f\|_\infty$, where

$$\|f\|_\infty \equiv \sup\{|f(x)| : a \leq x \leq b\} = \max\{|f(x)| : a \leq x \leq b\}.$$

Note: In the metric space $(C[a, b], d)$ with $d(f, g) = \|f - g\|_\infty$, convergence of sequences of “points”, $f_n \xrightarrow{n \rightarrow \infty} f$, implies $f_n \xrightarrow[n \rightarrow \infty]{\text{unif}} f \in C[a, b]$.

Example

In the metric space $C([a, b])$, with distance given by the sup-norm,

$$d(f, g) = \|f - g\|_\infty = \sup\{|f(x) - g(x)| : a \leq x \leq b\},$$

let


$$E = \{f \in C([a, b]) : 0 < f(x) < 1, \forall x \in [a, b]\}.$$

What are the interior E° and closure \bar{E} of the set E ?

Topology of metric spaces

Example (Cubic balls in $C[0, 1]$)

Suppose $f, g \in E \subset C[0, 1]$, $d(f, g) = \|f - g\|_\infty$.




The vertical **black line** indicates the distance between f and g .

Topology of metric spaces

Example (Cubic balls in $C[0, 1]$)

Suppose $f, g \in E \subset C[0, 1]$, $d(f, g) = \|f - g\|_\infty$.




The vertical **black line** indicates the distance between f and g .

Topology of metric spaces

Example (Cubic balls in $C[0, 1]$)

Suppose $f, g \in E \subset C[0, 1]$, $d(f, g) = \|f - g\|_\infty$.




The vertical **black line** indicates the distance between f and g .

Topology of metric spaces

Example (Cubic balls in $C[0, 1]$)

Suppose $f, g \in E \subset C[0, 1]$, $d(f, g) = \|f - g\|_\infty$.



The vertical **black line** indicates the distance between f and g .

Topology of metric spaces

Example (E° and \bar{E} for $E = \{f \in C([a, b]) : 0 < f(x) < 1 \quad \forall x \in [a, b]\}$)

If $f \in C([a, b])$ then f is continuous on $[a, b]$ so, by the Extreme Value Theorem, f attains its minimum and maximum values on $[a, b]$, say at $u, v \in [a, b]$. Since $f(x) \in (0, 1)$ for all $x \in [a, b]$, the extreme values, $f(u)$ and $f(v)$, must also be in the open interval $(0, 1)$. Therefore,

$$0 < f(u) \leq f(x) \leq f(v) < 1, \quad \forall x \in [a, b]. \quad (\heartsuit)$$

Thus, the range of f is $[f(u), f(v)] \subset (0, 1)$. There is a **gap** (a finite interval) that “insulates” f from the extreme values (0 and 1).

Let $\varepsilon = \min\{f(u), 1 - f(v)\}$. Then $\varepsilon > 0$. Now consider $g \in B_\varepsilon(f) \subset C([a, b])$.

$$\begin{aligned} \text{Then:} \quad & \|g - f\|_\infty < \varepsilon \\ & \implies \max\{|g(x) - f(x)| : 0 \leq x \leq 1\} < \varepsilon \\ & \implies |g(x) - f(x)| < \varepsilon \quad \forall x \in [0, 1] \\ & \implies -\varepsilon < g(x) - f(x) < \varepsilon \quad \forall x \in [0, 1] \\ & \implies f(x) - \varepsilon < g(x) < f(x) + \varepsilon \quad \forall x \in [0, 1] \end{aligned}$$

Topology of metric spaces

Example (E° and \bar{E} for $E = \{f \in C([a, b]) : 0 < f(x) < 1 \ \forall x \in [a, b]\}$)

Now using (\heartsuit), we have

$$0 \leq f(u) - \varepsilon \leq f(x) - \varepsilon < g(x) < f(x) + \varepsilon \leq f(v) + \varepsilon \leq 1, \quad \forall x \in [a, b],$$

from which we conclude that $g \in E$.

Since g was an arbitrary “point” in $B_\varepsilon(f)$, it follows that $B_\varepsilon(f) \subseteq E$, so any $f \in E$ is an interior point, so E is **open** and $E^\circ = E$.

What about \bar{E} , the closure of E ?

We will show that the closure of E is the set

$$F = \{f \in C([a, b]) : 0 \leq f(x) \leq 1, \ \forall x \in [a, b]\}. \quad (\spadesuit)$$

We will show $\bar{E} \subseteq F$ and then $F \subseteq \bar{E}$.

Topology of metric spaces

Example (E° and \bar{E} for $E = \{f \in C([a, b]) : 0 < f(x) < 1 \quad \forall x \in [a, b]\}$)

($\bar{E} \subseteq F$) First, consider any sequence $(f_n)_{n \in \mathbb{N}}$ with $f_n \in E$, $\forall n \in \mathbb{N}$, and suppose $f_n \xrightarrow{n \rightarrow \infty} f$ in the sup-norm, i.e., $f_n \xrightarrow[n \rightarrow \infty]{\text{unif}} f$. Since uniform convergence implies pointwise convergence, and $f_n(x) \in (0, 1)$, we must have $0 \leq f(x) \leq 1$, $\forall x \in \mathbb{N}$, and since $f_n \xrightarrow[n \rightarrow \infty]{\text{unif}} f$, we know $f \in C([a, b])$, so $f \in F$. Since f is a limit point of E and any limit point of E must lie in \bar{E} , we must have $\bar{E} \subseteq F$.

($F \subseteq \bar{E}$) Suppose $f \in F$, and define the sequence $f_n \in E$ by

$$f_n(x) = \begin{cases} 1 - \frac{1}{n}, & \text{if } f(x) > 1 - \frac{1}{n}, \\ f(x), & \text{if } \frac{1}{n} \leq f(x) \leq 1 - \frac{1}{n}, \\ \frac{1}{n}, & \text{if } f(x) < \frac{1}{n}. \end{cases}$$

By construction, $f_n \xrightarrow{n \rightarrow \infty} f$, and so f is a limit point of E so (by the [theorem about closures](#) again) $F \subseteq \bar{E}$. *(Note: $f_n \rightarrow f$ is illustrated in the next few slides.)*

Therefore, $F = \bar{E}$. □

Illustration of $f_n \rightarrow f$ for $f_n \in E$, $f \notin E$, $f \in F$




Illustration of $f_n \rightarrow f$ for $f_n \in E$, $f \notin E$, $f \in F$




Illustration of $f_n \rightarrow f$ for $f_n \in E$, $f \notin E$, $f \in F$




Illustration of $f_n \rightarrow f$ for $f_n \in E$, $f \notin E$, $f \in F$




Illustration of $f_n \rightarrow f$ for $f_n \in E$, $f \notin E$, $f \in F$




Illustration of $f_n \rightarrow f$ for $f_n \in E$, $f \notin E$, $f \in F$




Illustration of $f_n \rightarrow f$ for $f_n \in E$, $f \notin E$, $f \in F$




Illustration of $f_n \rightarrow f$ for $f_n \in E$, $f \notin E$, $f \in F$





Illustration of $f_n \rightarrow f$ for $f_n \in E$, $f \notin E$, $f \in F$





Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 27
Metric Spaces V
Friday 21 March 2025

Announcements

- *Last time:* interior and closure
- I added **some graphs** illustrating $f_n \rightarrow f$ in the construction that proved $F \subseteq \overline{E}$ at the end of the discussion of E° and \overline{E} for $E \subset C[a, b]$.

Boundary

Definition (Boundary Point)

If $E \subseteq X$ in a metric space (X, d) , then x is a **boundary point** of E if every neighbourhood of x contains at least one point of E and at least one point not in E , *i.e.*,

$$\begin{aligned} \forall \varepsilon > 0 \quad & B_\varepsilon(x) \cap E \neq \emptyset \\ & \wedge \quad B_\varepsilon(x) \cap (X \setminus E) \neq \emptyset. \end{aligned}$$

Definition (Boundary)

If $E \subseteq X$ then the **boundary** of E , denoted ∂E , is the set of all boundary points of E .

Topology of metric spaces

Example (Properties of boundary of E in a metric space (X, d))

For any $E \subset X$

- $\partial E = \overline{E} \setminus E^\circ$;
- ∂E is a closed set;
- E is closed if and only if $\partial E \subseteq E$.

Excellent exercises...

Topology of metric spaces

Example $(E^\circ, \bar{E}, \partial E$ for $E = (0, 1)^\infty \subset \ell^\infty$)

In the metric space ℓ^∞ , i.e., bounded sequences (x_n) with distance given by the sup-norm,

$$d(x, y) = \|x - y\|_\infty = \sup\{|x_n - y_n| : n \in \mathbb{N}\},$$

let

$$E = \{(x_n) \in \ell^\infty : 0 < x_n < 1, \forall n \in \mathbb{N}\}.$$

What are the interior E° , closure \bar{E} , and boundary ∂E of the set E ?

Please do poll: Metric spaces: ℓ^∞

Consider the following points in ℓ^∞ .

Are they in E ? Are they in E° ? Are they in \bar{E} ? Are they in ∂E ?

- $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \dots)$
- $(0, 0, 0, \dots)$
- $(0, 1, 0, 1, \dots)$
- $((-1)^n)_{n \in \mathbb{N}}$
- $(\frac{1}{n})_{n \in \mathbb{N}}$
- $(\frac{1}{n+1})_{n \in \mathbb{N}}$

Continuity in metric spaces

We have previously considered continuous functions $f : \mathbb{R} \rightarrow \mathbb{R}$.

What does it mean for a function to map one metric space to another *continuously*?

Definition (Continuous function)

Suppose (\mathcal{M}, d) and (\mathcal{N}, ρ) are metric spaces, $f : \mathcal{M} \rightarrow \mathcal{N}$, and $x \in \mathcal{M}$. The function f is *continuous* at x if:

$$\forall \varepsilon > 0, \exists \delta > 0 \text{ } \vdash \text{ } (y \in \mathcal{M} \wedge d(x, y) < \delta) \implies \rho(f(x), f(y)) < \varepsilon.$$

If f is continuous at every $x \in \mathcal{M}$, we say that f is *continuous on \mathcal{M}* .

- Continuity is determined point-by-point, but it is not enough to know f at a point; we must know how f behaves in a neighborhood of the point.
- While (\mathcal{M}, d) and (\mathcal{N}, ρ) can be any metric spaces, the most common situations are $(\mathcal{N}, \rho) = (\mathbb{R}, \text{standard})$ (so $f : \mathcal{M} \rightarrow \mathbb{R}$) and $(\mathcal{M}, d) = (\mathcal{N}, \rho)$ (so $f : \mathcal{M} \rightarrow \mathcal{M}$).
- If \mathcal{M} is a function space (e.g., $C[a, b]$) then $f : \mathcal{M} \rightarrow \mathbb{R}$ is often called a *functional* and $f : \mathcal{M} \rightarrow \mathcal{M}$ is often called an *operator*.

Continuity in metric spaces

It is usually helpful to rephrase the **definition** of continuity more geometrically in terms of balls. If we write the ball of radius δ in the distance d on \mathcal{M} as

$$B_\delta^d(x) = \{y \in \mathcal{M} : d(x, y) < \delta\},$$

then

$$y \in \mathcal{M} \wedge d(x, y) < \delta \iff y \in B_\delta^d(x).$$

To rephrase the second part of the **definition**, for any subset $E \subseteq \mathcal{M}$, we write the **image** of E by f as

$$f(E) = \{f(x) : x \in E\} \subseteq \mathcal{N}.$$

Then, we can write " $f(x) \in B, \forall x \in A$ " as " $f(A) \subseteq B$ ", so the **definition** of continuity can be expressed concisely as

$$\forall \varepsilon > 0, \exists \delta > 0 \quad \} \quad f(B_\delta^d(x)) \subseteq B_\varepsilon^p(f(x)).$$

So a function is continuous at $x \in \mathcal{M}$ if any ball about $f(x) \in \mathcal{N}$ is the image of a ball about $x \in \mathcal{M}$.



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 28
Metric Spaces VI
Monday 24 March 2025

Announcements

Assignment 5 is posted on the course web site.

Last time...

- boundary
- $(0, 1)^\infty$ in ℓ^∞
- **continuous functions** from one metric space to another
- continuity expressed with **balls**

Continuity in metric spaces

As in \mathbb{R} , in a general metric space we can express continuity using sequences.

Theorem (Continuity via sequences)

Let (\mathcal{M}, d) and (\mathcal{N}, ρ) be metric spaces, and suppose $f : \mathcal{M} \rightarrow \mathcal{N}$. Then f is continuous at $x \in \mathcal{M}$ if and only if for any sequence $(x_n)_{n \in \mathbb{N}} \subseteq \mathcal{M}$

$$x_n \xrightarrow{n \rightarrow \infty} x \implies f(x_n) \xrightarrow{n \rightarrow \infty} f(x).$$

Thus, f is continuous on \mathcal{M} iff, for every convergent sequence (x_n) ,

$$\lim_{n \rightarrow \infty} f(x_n) = f\left(\lim_{n \rightarrow \infty} x_n\right).$$

Proof.

(\implies) Suppose f is continuous at $x \in \mathcal{M}$, and $x_n \xrightarrow{n \rightarrow \infty} x \in \mathcal{M}$. By [definition](#), given any $\varepsilon > 0$, $\exists \delta > 0$ such that

$$y \in \mathcal{M} \wedge d(x, y) < \delta \implies \rho(f(x), f(y)) < \varepsilon. \quad (*)$$

But $x_n \xrightarrow{n \rightarrow \infty} x$ implies $\exists N \in \mathbb{N} \ \forall n \geq N, d(x, x_n) < \delta$. So $(*)$ implies $\rho(f(x), f(x_n)) < \varepsilon$, i.e., $f(x_n) \xrightarrow{n \rightarrow \infty} f(x)$.

... continued ...

Continuity in metric spaces

Proof of continuity via sequences (continued).

(\Leftarrow) Suppose that $x_n \xrightarrow{n \rightarrow \infty} x \implies f(x_n) \xrightarrow{n \rightarrow \infty} f(x)$ but f is not continuous at x . Then, inverting the definition, $\exists \varepsilon_0 > 0$ such that

for any $\delta > 0$ we can find $y = y(\delta) \in \mathcal{M}$
with $d(x, y) < \delta$ and $\rho(f(x), f(y)) \geq \varepsilon_0$.

In particular, for $\delta = \frac{1}{n}$, where $n \in \mathbb{N}$, $\exists y_n \in \mathcal{M}$ with

$$d(x, y_n) < \frac{1}{n} \quad \text{and} \quad \rho(f(x), f(y_n)) \geq \varepsilon_0.$$

This sequence (y_n) converges to x , so by hypothesis $f(y_n) \xrightarrow{n \rightarrow \infty} f(x)$. But this contradicts $\rho(f(y_n), f(x)) \geq \varepsilon_0 \implies \Leftarrow$ Hence f is continuous at x . \square

Continuity in metric spaces

Example (Continuity in $(\mathbb{R}, \text{standard})$ (refresher exercises))

1 Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$, f is continuous at all $x \in \mathbb{R}$, and $f(q) = 0$ for all $q \in \mathbb{Q}$. Prove that $f(x) = 0, \forall x \in \mathbb{R}$.

2 Let $f : \mathbb{R} \rightarrow \mathbb{R}$,

$$f(x) = \cos \frac{1}{x}, \quad \forall x \neq 0, \quad f(0) = a.$$

Show that no matter how we choose the value of $a \in \mathbb{R}$, f is never continuous at $x = 0$.

Continuity in metric spaces

Another equivalent characterization of **continuity** depends on the notion of:

Definition (Preimage or inverse image of a set)

Let $f : \mathcal{M} \rightarrow \mathcal{N}$ and $A \subseteq \mathcal{N}$. The **preimage** or **inverse image** of a A with respect to f is the set

$$f^{-1}(A) = \{x \in \mathcal{M} : f(x) \in A\}.$$

Note: The function f in this definition does not need to be invertible. There may be many points in \mathcal{M} that f maps to a single point in \mathcal{N} .

Continuity in metric spaces

Theorem (Continuity via inverse images)

Let $f : \mathcal{M} \rightarrow \mathcal{N}$. The following are equivalent:

- 1 f is continuous on \mathcal{M} .
- 2 If $U \subset \mathcal{N}$ is an open set in \mathcal{N} , then $f^{-1}(U)$ is an open set in \mathcal{M} .
- 3 If $F \subset \mathcal{N}$ is a closed set in \mathcal{N} , then $f^{-1}(F)$ is a closed set in \mathcal{M} .

Note: f continuous does not imply that f maps open sets to open sets. $f(U)$ need not be open if U is open, and $f(F)$ need not be closed if F is closed.

Example

Consider $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = 1 \forall x$.

$U = (0, 1)$ is an open set, but $f(U) = \{1\}$ is not open.

Consider $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = e^x \forall x$.

$F = \mathbb{R}$ is a closed set, but $f(F) = (0, \infty)$ is not closed.

Note: There is no mention of the metrics in the theorem above. If there is no metric on a space then we define continuity via inverse images.

Continuity in metric spaces

Proof of continuity via inverse images.

(1 \implies 3) Let $E \subset \mathcal{N}$ be any closed set, and take any sequence $(x_n)_{n \in \mathbb{N}}$ with $x_n \in f^{-1}(E)$, $\forall n \in \mathbb{N}$, and $x_n \rightarrow x$ in (\mathcal{M}, d) . Since f is continuous, it follows that $f(x_n) \xrightarrow{n \rightarrow \infty} f(x)$ in (\mathcal{N}, ρ) . Since E is closed, $f(x) \in E$, which is the same as saying $x \in f^{-1}(E)$. Therefore, $f^{-1}(E)$ is closed, so 3 holds.

(3 \implies 2) Let $V \subseteq \mathcal{N}$ be any open set. Then V^c is closed in (\mathcal{N}, ρ) . But $f^{-1}(V^c) = \{x \in \mathcal{M} : f(x) \in V^c\} = \{x \in \mathcal{M} \mid f(x) \notin V\} = (f^{-1}(V))^c$, so by 3 $f^{-1}(V)$ is open, i.e., 2 holds.

(2 \implies 1) Let $\varepsilon > 0$ and $x \in \mathcal{M}$ be given. Since $V = B_\varepsilon^\rho(f(x))$ is open in (\mathcal{N}, ρ) , $f^{-1}(V)$ is open in (\mathcal{M}, d) . Therefore, x is an interior point of $f^{-1}(V)$, so $\exists \delta > 0$ for which $B_\delta^d(x) \subseteq f^{-1}(V)$, or, $f(B_\delta^d(x)) \subseteq V = B_\varepsilon^\rho(f(x))$. Hence, we conclude that f is continuous at any $x \in \mathcal{M}$, i.e., 1 holds. \square

Continuity in metric spaces

Example (Using inverse images to prove continuity)

Let's revisit the first [refresher exercise](#), but prove it using the [inverse image of closed sets](#) (3) characterization of continuity.

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous at all $x \in \mathbb{R}$, and $f(q) = 0 \forall q \in \mathbb{Q}$.

Which closed set in $\mathcal{N} = \mathbb{R}$ should we focus on?

The range of f contains 0, and that's all we know about the range of f . So consider $F = \{0\}$.

F is closed, and $f^{-1}(F)$ contains \mathbb{Q} , since $f(q) = 0 \forall q \in \mathbb{Q}$. But f is [continuous](#). So $f^{-1}(F)$ is closed. Therefore, $f^{-1}(F)$ contains all the accumulation points of \mathbb{Q} , i.e., $f^{-1}(F)$ contains all of \mathbb{R} .

But $\mathbb{R} = f^{-1}(F) \implies f(\mathbb{R}) = f(f^{-1}(F)) = F = \{0\}$, i.e., $f(x) = 0$ for all $x \in \mathbb{R}$. □

Continuity in metric spaces

Example (Good exercises involving the discrete metric)

- 1 Consider $f : \mathcal{M} \rightarrow \mathcal{N}$ with domain $(\mathcal{M}, \text{discrete})$, and \mathcal{N} any metric space (\mathcal{N}, ρ) . Show that any such f is continuous.
- 2 Now suppose $f : \mathcal{M} \rightarrow \mathcal{N}$ but the range is $(\mathcal{N}, \text{discrete})$, and (\mathcal{M}, d) is a metric space that is not discrete (where “discrete” means $A \subset \mathcal{M}$ is both open and closed iff $A = \mathcal{M}$ or $A = \emptyset$). Show that f is a constant function.

Continuity in metric spaces

Example ($f : [0, 1] \rightarrow \mathbb{R}$)

Let $f : [0, 1] \rightarrow \mathbb{R}$, with the usual metric on $\mathcal{M} = \mathbb{R} = \mathcal{N}$, and $f(x) = x^2$ for all $x \in [0, 1]$. Then $V = (\frac{1}{4}, 2)$ is open in \mathbb{R} , and

$$f^{-1}(V) = \left\{ x \in [0, 1] : x^2 \in \left(\frac{1}{4}, 2 \right) \right\} = \left(\frac{1}{2}, 1 \right],$$

which—as a subset of \mathbb{R} with the usual metric—is neither open nor closed! *What went wrong?*

Nothing is wrong! The domain is the metric space $[0, 1]$, not \mathbb{R} , and the set $(\frac{1}{2}, 1]$ is open in $[0, 1]$. Since the domain space is $[0, 1]$, no subset can contain points that are not in $[0, 1]$. Balls in $[0, 1]$ mean the intersection of balls in \mathbb{R} with $[0, 1]$. This is called the *relative topology* (or *subset topology* or *induced topology*).



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 29
Metric Spaces VII
Wednesday 26 March 2025

Announcements

Assignment 5 is posted on the course web site.

Last time...

- continuity via sequences
- continuity via inverse images

Continuity in metric spaces

Definition (Open or closed relative to $A \subset \mathcal{M}$)

- 1 $U \subset A$ is **open relative to A** if $U = V \cap A$ for an open set V of \mathcal{M} .
- 2 $F \subset A$ is **closed relative to A** if $F = E \cap A$ for a closed set E of \mathcal{M} .

So the set $U = (\frac{1}{2}, 1]$ is open relative to $A = [0, 1]$ in \mathbb{R} , because (for example) $U = (\frac{1}{2}, 2) \cap A$, with $(\frac{1}{2}, 2)$ an open set in \mathbb{R} .

To distinguish a function $f : \mathcal{M} \rightarrow \mathcal{N}$ with domain $A \subset \mathcal{M}$ from a function defined on all of \mathcal{M} , we use the notation $f|_A$ for the restricted function (with domain A) that agrees with $f(x)$ for $x \in A$.

Example (Restriction of $f : \mathbb{R} \rightarrow \mathbb{R}$ to $A \subset \mathbb{R}$)

$$\text{Let } f(x) = \begin{cases} -x, & x \leq 0, \\ 1, & x > 0, \end{cases} \quad \text{and } A = (-\infty, 0].$$

Then $f|_A = -x$ is continuous on A , and, in particular, $f|_A$ is continuous on the boundary of A , i.e., $f|_A$ is continuous at $x = 0$.

Continuity in metric spaces

Continuity in subspaces of metric spaces:

In general, the equivalence of the **inverse image definition** and the **sequence definition** of **continuity** applies for functions $f|_A$ that are restricted to the domain $A \subset \mathcal{M}$ provided we define “open” and “closed” in the relative topology on A , and restrict to sequences $(x_n)_{n \in \mathbb{N}}$ with $x_n \in A \forall n \in \mathbb{N}$.

Continuity of operators on metric spaces of functions:

Remember that continuity of an operator $T : \mathcal{M} \rightarrow \mathcal{N}$ is distinct from the question of whether elements (*i.e.*, functions) in \mathcal{M} and \mathcal{N} are continuous.

Continuity in metric spaces

Example (Is the integral operator continuous?)

Consider the operator $T : C[a, b] \rightarrow C[a, b]$ with $T(f)$ defined by

$$T(f)(x) = \int_a^x f(x) dx.$$

Is T continuous?

Note: If $g = T(f)$, then g is continuous by the “integrals are continuous” theorem or the FFTC (since f is continuous).

The question is not whether such a g is continuous. It is whether T is continuous. Phrased in terms of **sequences**, the question is:

If $f_n \rightarrow f \in C[a, b]$ does it follow that $T(f_n) \rightarrow T(f) \in C[a, b]$?

Does the answer depend the metric?

Continuity in metric spaces

Example (Is the integral operator continuous?)

The answer is yes, for the sup norm on $C[a, b]$, $\|f\|_\infty = \sup_{x \in [a, b]} |f(x)|$.

Proof: For any $f_n, f \in C[a, b]$, and any $x \in [a, b]$, we have

$$\begin{aligned}
 |T(f_n)(x) - T(f)(x)| &= \left| \int_a^x (f_n(t) - f(t)) dt \right| && \text{now apply the triangle inequality} \\
 &\leq \int_a^x |f_n(t) - f(t)| dt \\
 &\leq \int_a^x \sup_{t \in [a, b]} |f_n(t) - f(t)| dt && (\clubsuit)
 \end{aligned}$$

If $f_n \xrightarrow{n \rightarrow \infty} f$ wrt $\|\cdot\|_\infty$, i.e., $f_n \xrightarrow[\text{unif}]{n \rightarrow \infty} f$, then $\forall \varepsilon > 0 \exists N \in \mathbb{N} \} \forall n \geq N$

$$\|f_n - f\|_\infty = \sup_{t \in [a, b]} |f_n(t) - f(t)| < \varepsilon. \quad (\spadesuit)$$

... continued ...

Continuity in metric spaces

Example (Is the integral operator continuous?)

Inserting (♠) in (♣), we have, for any $x \in [a, b]$,

$$|T(f_n)(x) - T(f)(x)| < \int_a^x \varepsilon dt = \varepsilon(x - a) \leq \varepsilon(b - a)$$

$$\therefore \|T(f_n) - T(f)\|_\infty = \sup_{x \in [a, b]} |T(f_n)(x) - T(f)(x)| \leq (b - a)\varepsilon.$$

Now, recognizing that in (♠) we can replace ε with $\frac{\varepsilon}{2(b-a)}$, we can conclude that $\forall \varepsilon > 0 \exists N \in \mathbb{N} \} \forall n \geq N, \|T(f_n) - T(f)\|_\infty < \varepsilon$, i.e., $T(f_n) \xrightarrow[\text{unif}]{n \rightarrow \infty} T(f)$.

Thus, the **integral operator** (T) is **continuous** on the metric space $C[a, b]$ with norm $\|\cdot\|_\infty$. □

Completeness

Completeness of metric spaces

The concept of a Cauchy sequence generalizes to any metric space.

Definition (Cauchy sequence in a metric space)

In a metric space (\mathcal{M}, d) , a sequence $(x_n)_{n \in \mathbb{N}}$ is called a **Cauchy sequence** iff

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \quad \vdash \quad m, n \geq N \implies d(x_n, x_m) < \varepsilon.$$

Theorem (Convergent implies Cauchy)

In a metric space (\mathcal{M}, d) , if (x_n) converges then (x_n) is a Cauchy sequence.

Proof.

Given $x_n \rightarrow x$, for any $\varepsilon > 0$, we can find $N \in \mathbb{N}$ such that $\forall n \geq N$, $d(x_n, x) < \frac{\varepsilon}{2}$. Suppose both $m \geq N$ and $n \geq N$. Then

$$d(x_n, x_m) \leq d(x_n, x) + d(x, x_m) \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

So (x_n) is Cauchy. □

Completeness of metric spaces

In $(\mathbb{R}, \text{standard})$, in addition to $\text{convergent} \implies \text{Cauchy}$ the converse $\text{Cauchy} \implies \text{convergent}$ is also true. *This converse does not hold in general in every metric space.* If it does hold, then we say the metric space is complete.

Definition (Complete metric space)

A metric space (\mathcal{M}, d) is said to be **complete** iff every Cauchy sequence in \mathcal{M} converges (to a point in \mathcal{M}).

Example (\mathbb{Q})

Let $q_n = \sum_{k=0}^n \frac{1}{k!}$. Then $q_n \in \mathbb{Q}$ for all n . But $q_n \rightarrow e = \sum_{k=0}^{\infty} \frac{1}{k!}$, and we showed $e \notin \mathbb{Q}$. So \mathbb{Q} is not complete.

In \mathbb{R} , the existence of least upper bounds is equivalent to Cauchy sequences always converging. So $(\mathbb{R}, \text{standard})$ is complete. But \mathbb{R} has much more structure. It is a field and it has an order. In fact, \mathbb{R} *is the unique, complete, ordered, field.*

Completeness of metric spaces

Example (Euclidean n -space)

- \mathbb{R}^n with the Euclidean norm is complete.

Proof: Let $(x_k)_{k \in \mathbb{N}}$ be a Cauchy sequence in $(\mathbb{R}^n, \text{Euclidean})$, say $x_k = (x_{k,1}, x_{k,2}, \dots, x_{k,n})$. For each fixed $j = 1, 2, \dots, n$, the sequence of real numbers $(x_{k,j})_{k \in \mathbb{N}}$ is a Cauchy sequence in \mathbb{R} , and hence converges in \mathbb{R} , say $x_{k,j} \xrightarrow{k \rightarrow \infty} \hat{x}_j \in \mathbb{R}$, for each $j = 1, 2, \dots, n$. Define $\hat{x} = (\hat{x}_1, \hat{x}_2, \dots, \hat{x}_n) \in \mathbb{R}^n$. Since each component of the vector converges, it follows (from A5 Q1) that $x_k \xrightarrow{k \rightarrow \infty} \hat{x}$. \square

Note: $(\mathbb{R}^n, \text{Euclidean})$ is an example of a **Hilbert Space**, an inner product space that is complete wrt the induced metric.

- \mathbb{R}^n is also complete wrt the Taxicab ($\|\cdot\|_1$) and Maximum ($\|\cdot\|_\infty$) norms (*check!*). In fact, for any $p \geq 1$, \mathbb{R}^n with norm $\|\cdot\|_p$ is complete wrt to the metric $d(x, y) = \|x - y\|_p$ induced by the norm.

Any complete, normed, vector space is said to be a **Banach Space**.

Poll

- Go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Metric spaces: completeness**
- .

Completeness of metric spaces

Example (More examples related to completeness)

- For any interval $I \subset \mathbb{R}$, the space $C_b(I)$ of bounded continuous functions, with distance determined by the sup-norm, is a complete metric space. *A good exercise.*

Hint: If (f_n) is a Cauchy sequence in $C_b(I)$ with the sup norm, what does that imply about sequences $(f_n(x))$ in \mathbb{R} for $x \in I$?

- Consider the subspace of $C[0, 1]$ consisting of polynomials. We call this $P[0, 1]$. *Is $P[0, 1]$ complete?* Can we construct a sequence of polynomials that does not converge in $P[0, 1]$? The sequence

$$P_n(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!}$$

converges as $n \rightarrow \infty$ (uniformly, *i.e.*, in the sup norm) to e^x , which is not a polynomial $\implies P[0, 1]$ is not complete.